HILGARDIA

JOURNAL OF AGRICULTURAL SCIENCE PUBLISHED BY
THE CALIFORNIA AGRICULTURAL EXPERIMENT STATION

Volume 37, Number 2 • December, 1965

Anatomy and Cytology of *Vitis* Phloem

Katherine Esau

UNIVERSITY OF CALIFORNIA DIVISION OF AGRICULTURAL SCIENCES

CONTENTS

Introduction	17
Material and methods	18
Anatomy of the Bark of <i>Vitis</i> species	19
Periderm	19
Phloem	21
Tylosoids in nonconducting phloem	25
Reactivation of phloem	27
Cellular differentiation in the root tip	27
Meristematic and cortical cells	27
Phloem cells	33
Secondary phloem in the stem	39
Contents of sieve elements	39
Companion cells	47
Sieve plates and lateral sieve areas	55
Parenchyma cells and fibers	59
Discussion	61
Summary	69
Acknowledgments	70
Literature cited	71

THE AUTHOR:

Katherine Esau is Professor of Botany, Emeritus, Davis.

Anatomy and Cytology of *Vitis* Phloem^{1,2}

INTRODUCTION

AN INTENSIVE investigation of the phloem of *Vitis vinifera* L., the grapevine, was previously reported in this Journal (Esau, 1948a).³ That study was centered on the development of the tissue, especially with reference to seasonal changes. It served as a background for the subsequent study of the effects of Pierce's disease virus on grapevine (Esau, 1948b). In connection with the earlier study, samples of bark were collected from thirty species of *Vitis*, other than *V. vinifera*, growing in the experimental vineyard of the Department of Viticulture at Davis. The first part of the present paper reports on a survey of phloem of these species.

After an electron microscope was installed on the Davis campus, the ultrastructure of phloem tissue became one of the important research projects in the Department of Botany at Davis. *Vitis* phloem was included in the studies and a considerable amount of information on this tissue was obtained during 1960–1962. This information constitutes the major part of the present paper.

Methodology in submicroscopic research is still new and is changing rapidly. The ultrastructural data reported here are based on relatively early techniques. Nevertheless, these data expand the understanding of the structure and seasonal changes in the conducting

element, the sieve element, and further differentiate this element from the associated nucleate cells. The present study has confirmed the interpretation of some previously recognized features but has also introduced new problems.

Early ultrastructural investigations on plant material have emphasized meristematic or relatively undifferentiated parenchyma cells, especially those derived from root tips (see, for example, Whaley *et al.*, 1960), and the submicroscopic structure of such cells is relatively well understood. Interpretation of the fine structure of mature and highly specialized cells must therefore be made by comparison with the less-differentiated cells. Part of the present paper deals with the sieve element in the root tip, which was studied against the background of information on more or less meristematic parenchyma cells of the same parts of the root. These background observations are reported in full because they help to single out the specific characteristics of the sieve element.

The literature pertinent to the subject matter is not reviewed under a separate heading. Certain references are mentioned in connection with individual observations in the main text, but the comprehensive comparison of data obtained with those reported in the literature appears only in the discussion. The specific terminology concerning the phloem tissue is explained in the previous Hilgardia article on *Vitis* phloem (Esau, 1948a), and that serving for the description of ultrastructural details has been reviewed by Esau and Cheadle (1965).

¹ Submitted for publication May 5, 1965.

² Part of the ultrastructural study reported here was presented as an invitational paper at the Tenth International Botanical Congress, August 1964, in Edinburgh, Scotland, entitled "Aspects of differentiation at ultrastructural level."

³ See "Literature Cited" for citations referred to in the text by author and date.

MATERIALS AND METHODS

The collections of species from the Department of Viticulture vineyard at Davis included branches ("canes" and "arms") of various ages. A major collection was made on January 10 and 11, 1946, when the cambium and phloem were dormant. For determination of possible reactivation of previously active but now dormant phloem—a relatively uncommon phenomenon but characteristic of *Vitis vinifera*—bark samples of the same species were collected on June 18, 1947. The bark was slipping at this time, an indication of cambial activity.

Dormant material was killed in chrome-acetic-formalin solution, embedded in water-soluble stearate, sectioned on a sliding microtome, and stained with Bismark brown, iodine green, and lacmoid (often called resorcin blue), all according to Esau (1948a). Summer material was killed in the same mixture, sectioned on a freezing microtome, and stained as the dormant material. In both sets of slides the sections were mounted in light Karo syrup and the cover glasses were sealed with Canada balsam. The two stains, Bismark brown and iodine green, have retained their brilliance through the years, but the lacmoid has largely faded.

Some of the root tip material used for light-microscope study was killed in chrome-acetic-formalin solution, processed through paraffin, and stained with hematoxylin and safranin according to the schedule described by Esau (1944). Other root tips were studied in slides used as monitors in electron microscopy. These slides bore sections, cut 1/2- to 1-micron thick, of material embedded in methacrylate and stained with safranin and methyl violet (Esau, Cheadle, and Risley, 1962).

For electron microscopy, most samples were fixed in 4 per cent potassium permanganate (KMnO_4) for several

minutes at room temperature (23°C) and all were embedded in methacrylate. Some material was fixed in osmic acid. The fixations are indicated in the legends. In nine treatments the canes (1-year-old stems) were cut into 8-inch pieces in the field and placed for several hours in 0.25 M buffered or unbuffered sucrose solution before treatment for electron microscopy. Some of these samples were tested for plasmolyzability of sieve elements with positive results. In four additional treatments the canes were not pretreated with sucrose but were immediately cut under water into segments suitable for sliding microtome sectioning. In one additional treatment immersion in water was omitted.

In treatment for electron microscopy the bark was sectioned on a sliding microtome 40 to 60 microns thick. In some treatments the sections were further dissected in the bathing sucrose solution, then transferred to the killing solution; in others, the permanganate solution was applied to the freshly-exposed surface of a block being sectioned on the sliding microtome. The block was then washed with alcohol and sectioned further.

Material fixed with the same chemical showed no variations ascribable to differences in pretreatment. All sieve areas encountered contained darkly-stained material in their pores.

The root tips for electron microscopy were cut under a sucrose solution into 1-mm lengths beginning with the apex. Attempts were made to exclude the root-cap. Each piece was then quartered lengthwise before fixation in 4 per cent KMnO_4 for several minutes at room temperature.

Most of the material for electron microscopy was derived from *V. vinifera* var. Sultanina (Thompson Seedless), but some came from Black Monukka and Emperor varieties.

ANATOMY OF THE BARK OF VITIS SPECIES

Periderm

The previously mentioned detailed study of the phloem of *Vitis vinifera* (Esau, 1948a) served as the basis for comparing the phloem of the thirty other available species of this genus. Most of these species are listed in Bailey and Bailey's (1949) *Hortus Second* and their names (given below) are spelled according to that reference work. Species names marked with asterisks are not listed by Bailey and Bailey (1949) but were furnished, as spelled, by the Department of Viticulture, University of California, Davis.

The following species are represented in the slide collection that served for the comparative study of *Vitis* phloem: *amurensis*, *andersonii**, *arizonica*, *Bailleyana*, *Berlandieri*, *californica*, *candicans*, *Champinii*, *cinerea*, *Coignetiae*, *cordifolia* (*V. vulpina*), *coriacea* (*V. Shuttleworthii*), *Doaniana*, *flexuosa*, *Girdiana*, *Labrusca*, *Lincecumii*, *Longii*, *monticola*, *Munsoniana*, *Piasezkii* (var. *Pagnuccii*), *retordii**, *riparia* (*V. vulpina* of some), *rotundifolia* (*Muscadinia rotundifolia*), *rubra* (*V. palmata*), *rupestris*, *slavinii**, *Solonis* (*V. Longii*), *treleasii**, *vulpina* (*V. cordifolia*).

The bark (all stem tissues located outside the vascular cambium) of *Vitis vinifera* L. (Esau, 1948a) illustrates characteristics of this stem region prevalent in the family Vitaceae (or Ampelidaceae; Metcalfe and Chalk, 1950). The cortex is short-lived because the periderm is formed deep in the stem at the end of the first season of growth of the shoot (cane). According to Metcalfe and Chalk (1950), the phellogen arises in the pericycle and cuts off the pericycle and cortex. In *Vitis* no pericycle as such is present: the starch sheath (endodermis), that is the innermost layer of the cortex, abuts on the primary phloem. After the first sieve tubes

are obliterated, primary phloem fibers differentiate next to the starch sheath. Metcalfe and Chalk (1950) describe the pericycle of *V. vinifera* as containing isolated strands of fibers. This treatment obscures the developmental stages of the stem and presents incorrectly the origin of the first phellogen. The outermost extraxylary fiber strands in the stem of *Vitis* are associated with the vascular bundles and constitute fibers of the no-longer-functioning protophloem; the panels of the parenchyma cells alternating with the fibers are interfascicular regions (primary rays, Esau, 1948a; primary medullary rays, Metcalfe and Chalk, 1950). The first phellogen arises partly in the primary phloem (metaphloem) on the inner side of the protophloem fibers, and partly in the interfascicular parenchyma (Esau, 1948a). Thus, when the cane matures at the end of the first season the periderm cuts off as a unit (ringbark) the cortex, the primary phloem fibers, and some interfascicular parenchyma. In subsequent years phellogen forms in the secondary phloem and consequently successive layers of secondary phloem are shed (fig. 1,A). In secondary phloem the phellogen is derived from subdividing parenchymatic elements of the tissue: ray cells and axial parenchyma cells.

The place of origin of the first periderm in the different species was determined by examining the bark of canes at the end of the first season's growth (mature canes of the viticulturist). In all species, except *V. rotundifolia*, the tissue underlying the periderm was secondary phloem with nonfunctioning remnants of metaphloem; that is, the cortex and the primary phloem fibers were absent. In *V. rotundifolia* the periderm did not remove the cortex; it arose superficially. The same position was occupied by the periderm of older branches (arms; fig. 1,B). As a result of

Fig. 1. Transverse sections of bark. Sliding microtome sections. The phloem is dormant. A. *Vitis monticola*. Phloem contains one annual increment. Inner layer of periderm has cut off phloem increment of preceding year. Still-older phloem was shed earlier. Numbers 44 and 45 indicate that layers of late phloem (lp) were formed in 1944 and 1945. B. *Vitis rotundifolia*. Bark contains cortex and both primary and secondary phloem. Fibers at right are protophloem fibers. Periderm arose near surface of stem. Rays became dilated with increase in width of stem. (Both, X98.) Details: f, fiber; lp, late phloem; s, sieve element; v, vessel.

the shallow origin of the periderm in *V. rotundifolia* the bark of older stems in this species contains several years growth of phloem, remnants of primary phloem, within which fibers are the most conspicuous components (cells labeled *fibers* in fig. 1,B), and the periderm. The rays widen in their outer parts by dilatation growth.

In *Vitis* species other than *V. rotundifolia*, layers of phloem are cut off yearly by periderm arising in successively deeper tissues. The bark is shed as a ringbark. In *V. rotundifolia* the stem remains smooth because only a small amount of cork is sloughing off. No collections were made to determine whether still-older stems develop deeper periderm in this species. Presumably the first periderm persists for a long time because Bailey and Bailey (1949) characterize the bark of *V. rotundifolia* as not shedding.

In species shedding phloem during periderm formation the amount of secondary phloem accumulating beneath the periderm is variable (fig. 2), and phellogen occasionally originates at different depths around the same circumference of the stem. In such instances the outer bark surface is uneven (fig. 2,B). Some undulation of the bark surface may result also from the circumstance that the phellogen has a deeper origin in the rays than it does elsewhere (figs. 1,A, and 2,A).

Because of the somewhat variable depth of formation of the successive phellogens, secondary phloem may be represented by one or more annual increments. The phloem in figure 1,A, for example, contains only one increment and a small amount of another; that in figure 2,B, contains one and parts of a second in the thinnest region and two and part of a third in the thickest. This interpretation is based on the assumption that the layers of narrow cells labeled *lp* in figures, 1,A, and 2,B, constitute late phloem—that is, phloem produced at the end of seasonal growth.

In the absence of layers of conspicuously narrow cells distally from cambium (fig. 2,A,C) the age of the bark is difficult to determine. Judged by the appearance of the outermost sieve tubes in figures 2,A and C, the entire thickness of phloem in these samples may have been formed in one season. If so, these views would exemplify an extra-vigorous growth of phloem.

The variations in bark structure just discussed, other than those found in *V. rotundifolia*, can hardly be considered specific variations even though they are illustrated here by reference to different species; such variations probably could be found in any species if large enough samplings were made. In *V. vinifera* studied earlier, for example, the number of annual increments accumulated under the periderm varied from one to four, and the thickness of the bark varied from $\frac{1}{2}$ to 2 millimeters (Esau, 1948a).

Phloem

The species of *Vitis* examined have generally similar phloem structure, except that some minor deviations were found in *V. rotundifolia*. The structure is that identified by Metcalfe and Chalk (1950) as characteristic of the *Euvitis* section of *Vitis*. The axial system of the phloem occurs in discrete blocks separated from one another by wide rays. Within the blocks, tangential strata of sieve elements with associated nucleate cells alternate with tangential bands of fibers (figs. 1,A, and 2). In *V. rotundifolia* the fibers are sparse and do not form regular tangential bands (fig. 1,B). The absence of tangential strata of fibers is given by Metcalfe and Chalk (1950) as a distinguishing characteristic of the *Muscadinia* section of *Vitis*. The fibers are septate (fig. 3), a family characteristic (Metcalfe and Chalk, 1950). In the present study septae were found in all species, including *V. rotundifolia*. The fibers are concerned with starch storage, as are the parenchymatic members of the phloem.

Fig. 2. Transverse sections of bark showing variations in thickness of secondary phloem. Sliding microtome sections. The phloem is dormant. A. *Vitis riparia*. B. *V. Berlandieri*. Numbers 43, 44, and 45 indicate that layers of late phloem (lp) were formed in 1943, 1944, and 1945. C. *V. Labrusca*. D. *V. Coignetiae*. (All, X70.) Details: lp, late phloem; s, sieve element; v, vessel.

Fig. 3. Tangential longitudinal section of secondary phloem of dormant cane collected Jan. 31, 1961. From monitor slide. Osmium fixation. Ray cells at edges, right and left, axial system in the middle. Dark contents in cells, tannins. Sieve element (s) with four companion cells (numbered 1-4) and compound sieve plates (pt) covered with dormancy callose. (X440). Details: f, septate fiber; pa, parenchyma cell; pt, sieve plate; ry, ray cell; s, sieve element.

Fig. 4. Tylosoids in nonfunctioning sieve elements. Sliding microtome sections. A. *Vitis andersonii*. Radial section through functional phloem showing nonoccluded sieve element (s) and associated parenchyma cells (pa and tpa). B. *V. Labrusca*. Tangential section through nonconducting phloem with two sieve elements occluded with tylosoids. Some of these have secondary cell walls (sw). C. *V. Labrusca*. Tangential section through nonconducting phloem showing part of a sieve tube with sieve plate (sp). Tylosoids (t) occur to the left of the plate, which bears definitive callose (ca). The element to the right of the sieve plate is not occluded. (A, B, X440; C, X600.) Details: ca, callose; f, fiber; pa, parenchyma cell; s, sieve element lumen; sp, sieve plate; sw, secondary wall; t, tylosoid; tpa, tanniniferous parenchyma cell.

The sieve element is as long as the considerably elongated cambial fusiform initial, and has compound sieve plates in the oblique end walls. Sieve plates appear in sectional view in a tangential section of the tissue (fig. 3) and in surface view in a radial section (fig. 4,A). Each sieve element has several short companion cells (four are seen in figure 3). Some parenchyma cells and ray cells are densely filled with tannins (black in figs. 1–5); others contain starch and small amounts or none of tannins (fig. 33, A,B). According to Metcalfe and Chalk (1950), Ampelidaceae have solitary crystals, clustered crystals (druses), and needle-shaped (raphides) crystals. The three kinds of crystals were found in variable numbers in all species examined. The solitary rhombohedral crystals and the druses commonly occur in the marginal ray cells, whereas the raphides are found deeper in the rays, confined to enlarged idioblasts.

Tylosoids in nonconducting phloem

Occlusion of nonconducting sieve elements by outgrowths from contiguous parenchyma cells was observed in *Vitis vinifera* (Esau, 1948a). These outgrowths have been referred to in the literature as tyloses. "Tylosoids" is a more appropriate term for them because tyloses, as they occur in the xylem, grow through pits in secondary walls between parenchyma cells and tracheary elements, whereas tylosoids refers to cell outgrowths not related to pits.⁴ The occlusion of sieve elements occurs by more or less localized growth of adjacent parenchyma cells. In some instances tylosoid growth in phloem appears to be restricted to primary pit-field regions,

⁴ Although the wood anatomist uses the term "tylosoid" to denote "proliferation of a thin-walled epithelial cell into an intercellular canal" (Committee on Nomenclature, 1957), he also states that a tylosoid differs from a tylose in that it does not pass through a pit. This is the sense in which tylosoid is used here.

in others it involves a broad expanse of the cell wall. The connections between tylosoids and the originating cells (numbered regions in figure 5,A) are therefore variable in width but are seldom narrow. In the xylem, the corresponding connections between tyloses and the originating cells are narrow and relatively uniform because their diameter is determined by the size of the pit through which the tylose grows (Esau, 1948b, p. 476).

In many species of plants, nonfunctioning sieve elements (and the associated companion cells) are crushed completely by surrounding parenchyma cells. Such crushing of sieve elements has been observed in primary phloem of *V. vinifera* (Esau, 1948a) but in secondary phloem the obliteration of sieve elements occurs largely by development of tylosoids. Growth of parenchyma cells that brings about complete crushing of phloem cells, and the development of tylosoids, are related phenomena differing from one another only quantitatively.

All *Vitis* species surveyed in this study showed tylosoid development. In some samples the amount of nonconducting phloem was small, but at least a few sieve elements with tylosoids were found in each.

The development of tylosoids profoundly modifies the appearance of phloem tissue. The tissue assumes an aspect of a homogeneous parenchyma containing starch and tannins and interspersed with bands of fibers. Both kinds of parenchyma cells, those containing starch and those depositing tannins, are forming tylosoids (figs. 4,B, and 5). The sieve element is in contact with parenchyma cells along its radial (fig. 3), or tangential (fig. 4,A), or both kinds of walls. Tylosoids appear to have the tendency to develop in series, either in the radial plane (fig. 5,A) or in the tangential (fig. 5,C), sometimes from two opposite sides in the same cell (arrows in fig. 5,B). As a re-

Fig. 5. Tylosoids in nonfunctioning sieve elements. Sliding microtome sections. A,B. *Vitis Munsoniana*. Radial sections of nonconducting phloem. In A, tylosoids (t) are seen in continuity with the originating parenchyma cells; in B, tylosoids invaded sieve element from two opposite sides (arrows). C. *V. Labrusca*. Tangential section of nonconducting phloem. Tylosoids (t) derived from tanniferous cells. (A, B, X520; C, X600.) Details: f, fiber; fo, fold in cell wall formed by growth of tylosoid; n, nucleus; pa, parenchyma cell; pi, pits; s, sieve element lumen; sw, secondary wall in tylosoid; t, tylosoid; tpa, tanniferous parenchyma cell; numbers 1,2,3, indicate constrictions between tylosoids and the cells originating them.

sult of such symmetrical growth pattern, sections not passing through the cells originating the tylosoids display the latter as rows of superimposed short parenchyma cells more or less completely occluding the lumina of the sieve elements (fig. 4,B). The presence of sieve plates in contact with such cells (fig. 4,C) clearly reveals their nature. Sieve plates of occluded sieve elements may bear definitive callose (fig. 4,C) and the not-occluded parts of cell lumina may be filled with remains of protoplasmic contents (fig. 5,A; residual material, Esau, 1948a, p. 245).

Reactivation of phloem

The phenomenon of reactivation of phloem after winter dormancy is best documented for *Vitis vinifera* (see Esau, 1948a, pp. 253–257). Dormant sieve elements have massive amounts of callose on sieve areas of all degrees of specialization. During reactivation, callose diminishes in amount and sieve areas assume the same condition which they attained originally during differentiation from the meristematic state. Because reactivation has been recorded in only a few species of dicotyledons, it was desirable to determine whether it occurs in *Vitis* species other than *V. vinifera*. As mentioned in "Materials and Methods," the bark of various species intended for study of phloem reac-

tivation was collected in early summer of 1947 when cambial activity was well under way and the bark was slipping. Studies on *V. vinifera* (Esau, 1948a, pp. 257–259) have shown that at the time of slippage of bark the old phloem is reactivated and new phloem is differentiating. The contrast between reactivated and new sieve tubes is considerable because the new sieve tubes have a highly turgid appearance and their sieve plates are relatively thin. The somewhat thicker sieve plates of the reactivated elements have deeply-stained slime in the pores—a reliable indication of reactivation.

Evidence of partial or complete reactivation of the 1946 phloem was observed in all species but three. In *V. Doaniana* the old phloem had no dormancy callose and no slime strands in the sieve plates. The tissue appeared to be nonfunctioning. The new phloem also exhibited no strands in the sieve areas. In *V. treleasii* dormancy callose was unusually small in amount and thick pads of callose occurred in the new phloem. In *V. slavini* new growth was very scanty and old phloem appeared nonfunctional, as it was devoid of callose and slime. Apparently these samples of phloem were not representative of normal tissue in the three species. The phenomenon of reactivation is probably characteristic of *Vitis* as a genus.

CELLULAR DIFFERENTIATION IN THE ROOT TIP

Meristematic and cortical cells

Meristematic cells of root tips have served for the initial development of the concepts of subcellular structure of plant cells (see especially Whaley *et al.*, 1960). The basic features recorded for root tip cells in other species are found in the corresponding cells of *Vitis*.

Nuclei have the typical double-layered envelope (this feature is not represented in the figures) with pores (figs. 6,A, and 7,A,) which appear to

be rather evenly distributed when seen in surface views of nucleus (fig. 7, B and C). At higher magnifications (fig. 7, C) the pores show a central granule, a feature also observed by Wooding and Northcote (1965) in nuclei in *Acer pseudoplatanus* phloem. Extensions of the nuclear envelope, which are indistinguishable from endoplasmic reticulum membranes, are common (fig. 6,A). Permanganate fixation frequently fails to bring out the nucleolus (fig. 6,A) or gives a faint image of the body (fig. 8).

Fig. 6. Electron micrographs of cells from transactions of first mm of root tip (rootcap excluded). Potassium permanganate fixation. A. Procambial cells with no discernible vacuoles. B. Cortical cell with vacuoles (v). Details: d, dictyosome; er, endoplasmic reticulum; m, mitochondrion; n, nucleus; po, pore in nuclear envelope; v, vacuole; w, cell wall. (Magnification of this and following micrographs is indicated by a line, the value of which in microns is given by the μ sign preceded by a number.)

When better defined the nucleolus appears to contain vacuoles and to have no sharp boundary (fig. 7,A); that is, the nucleolus has no envelope of its own. Nuclei of meristematic and partly-differentiated cells may be more or less lobed (figs. 8,A and 12,A).

The endoplasmic reticulum is a relatively inconspicuous structural element of meristematic cells, but may be quite abundant in differentiating cells (compare figs. 6,A, 7,A, and 8 with figs. 9,A, 10,A, and 11). In permanganate-treated material it exhibits the common double-layered profiles. Bias cut of endoplasmic membranes may reveal their cisternoid form in both meristematic and differentiating cells (figs. 8,A, and 10,A). In some differentiating cells certain circular profiles suggest occurrence of tubular endoplasmic membranes or a dissociation of the membranes into vesicles (fig. 9,A, at **ve**). The other representative of membrane systems, the dictyosome (Golgi body), is a common component of meristematic and differentiating cells. In young phloem cells dictyosomes may be associated with numerous vesicles, a phenomenon that needs further study in view of the possible involvement of dictyosome vesicles in wall formation.

Mitochondria and plastids usually are difficult to distinguish from one another in meristematic cells. With advance in differentiation the development of numerous tubular or saccate membranes from the inner layer of the double-layered outer membrane serves to identify the mitochondria. Plastids remain poorly equipped with inner membranes but commonly produce starch (fig. 8,B). Indication of starch development—electron-transmitting regions mostly lacking starch as such—should not be confused with light regions in younger plastids and mitochondria (fig. 7,A). Whaley *et al.* (1960) have called attention to this phenomenon in mitochondria and have suggested that it may be an artifact in part.

In some meristematic cells the profiles of one or more mitochondria appear unusually long (figs. 6,A, and 8). These mitochondria have a striking resemblance with mitochondria in *Anthoceros* described by Manton (1961) as discs with thin centers bent into a bowl or saucer shape. They also remind one of mitochondria in the egg cell of *Pteridium* interpreted by Bell and Mühlethaler (1964) as having the shape of an umbo. To recognize the bowl (or umbo) shape, serial sections or sections cut parallel with the plane of the bowl are necessary. Illustrations of the latter type of sections are provided by the authors named above. In *Vitis*, evidence that mitochondria with the long profiles are not cylindrical but have an extended surface was found in views like that shown in figure 12,B. The apparently branched mitochondrion in this figure is most likely an incomplete section of a structure with an extended surface. Bell and Mühlethaler (1964) ascribe the umbo shape of *Pteridium* mitochondria to their origin from the nucleus, but Manton (1961) suggests that the bowl-shaped mitochondria are temporary configurations associated with mitochondrial growth and multiplication. The study of the large mitochondria in *Vitis* was not carried far enough to discuss their possible origin and significance.

Two kinds of vacuoles were observed in the differentiating cells—vacuoles free of tannin, and tanniniferous vacuoles; both kinds appeared to be of the type not related to endoplasmic membranes (see Esau, 1963, for review of the matter). The tannin-free vacuoles contained a thin precipitate, a common appearance in electron micrographs of plant cells (Whaley *et al.*, 1960). Manton (1962) states that commonplace materials such as soluble carbohydrates have the capacity to form vacuolar precipitates with potassium permanganate.

Tannin vacuoles containing phenolic compounds, or tannins, are widely dis-

Fig. 7. Electron micrographs of meristematic root tip cells illustrating nuclear details. Potassium permanganate fixation. A. Nucleus with pore in envelope. Nucleolus (nu) with vacuoles. B,C. Nuclear envelope seen from the surface at two magnifications. Details: er, endoplasmic reticulum; i, intercellular space; m, mitochondrion; nu, nucleolus; pl, plastid; po, pore in nuclear envelope; w, cell wall.

Fig. 8. Electron micrographs of meristematic cortical cells from root tip. Potassium permanganate fixation. A. Some mitochondria have extremely long profiles. They may be umbo or bowl shaped (fig. 12,B). B. Mitochondrion with long profile and plastids with starch grains. Details: arrow, branched plasmodesma with median nodule; d, dictyosome; er, endoplasmic reticulum; m, mitochondrion; n, nucleus; nu, nucleolus within nucleus; pl, plastid; po, pore in nuclear envelope; w, cell wall.

Fig. 9. Electron micrographs of cortical root tip cells. Potassium permanganate fixation. A. Cell with larger and smaller tannin vacuoles (v) and region with vesicles probably derived from endoplasmic reticulum. B. Cell with only few organelles and membrane systems and with a relatively dense hyaloplasm. C. Crystal cell. Sharply delimited portions of cytoplasm with dense hyaloplasm resembling that in B. Angular spaces where crystals (raphides) were present before fixation. Details: d, dictyosome; er, endoplasmic reticulum; m, mitochondrion; n, nucleus; pl, plastid; r, site of raphide crystal; v, vacuole; ve, vesicles; w, cell wall.

tributed in *Vitis* tissues (Schanderl, 1957) and occur in both cortex and vascular cylinder in the root. Tannin occurs throughout the vacuole, more or less dispersed (fig. 9,A), but usually also aggregates at vacuole periphery (fig. 11,A). In young vacuoles the peripheral accumulation appears extremely dense (fig. 10,B). The apparently solid vacuoles may be sections through such peripheral accumulations (fig. 10,B, arrow). At the ultrastructural level, Wardrop and Cronshaw (1962) relate the origin of phenolic substances in wood-ray parenchyma to dissolution of starch, and they describe accumulation of these substances in vesicles contained within chloroplasts or amyloplasts that later disintegrate and release the phenolic substances, presumably into vacuoles. On the basis of a light-microscope study of *Vitis* tissues, Schanderl (1957) relates tannin origin to plastid-like structures in callus and to "hollow bodies" in root endodermis. Only vacuolar position of tannin was observed in material used in the present study.

Many electron micrographs reveal no vacuoles in meristematic cells (fig. 6,A; see also fig. 2 in Whaley *et al.*, 1960). Light-microscope studies have led to the concept that presence of vacuoles is a general characteristic of meristematic cells. They are small and dispersed in smaller cells, large and conspicuous in larger cells (Zirkle, 1932). Variations in cells, differences in fixation, and thickness of sections may account for the discrepancy between the images produced by the two kinds of microscopes.

Root cortex is characterized by an early development of intercellular spaces. In material fixed with potassium permanganate, intercellular spaces in *Vitis* roots frequently contain a precipitate of unknown origin and composition (fig. 10,A). According to Genève (1964), intercellular spaces in radish root arise lysigenously but the debris of the degenerated cells are absorbed by contiguous cells. Lysigenous origin of

intercellular spaces was not recognized in *Vitis* roots.

Vitaceae (Ampelidaceae) are characterized by presence of raphide idioblasts (Metcalfe and Chalk, 1950). In leaves and in rays of secondary phloem of *Vitis* the raphide crystals occur in individual enlarged cells, the raphide sacs. In root cortex, however, the raphide idioblasts form continuous longitudinal series of cells that undergo considerable elongation after the crystals begin to form in them (fig. 13,C). According to Cheavin (1938), raphides occur in a mucilaginous vacuole. In material processed for electron microscopy the crystals are usually not preserved, but their sites within the vacuole remain discernible as electron-transmitting regions (figs. 9,C, and 11,B); these sites may have an angular shape in transections (r in fig. 9,C). Occasionally, crystals are preserved and appear as extremely opaque angular bodies in transections. Early stages of raphide idioblast development were not recognized with certainty. The hyaloplasm of raphide cells may appear dense (fig. 9,C). Cells with similar hyaloplasm and few organelles (fig. 9,B) might possibly be raphide idioblasts in early stages of differentiation. In some raphide cells most of the organelles and membrane systems appeared disorganized (fig. 9,C); in other, possibly younger, cells a considerable amount of endoplasmic reticulum was observed. The membranes were sometimes aggregated into coiled structures (fig. 11,B).

Phloem cells

Sieve elements of the protophloem in roots commonly are the first vascular cells to mature. They appear as clear cells having little stainable content and therefore they stand out sharply from associated cells having dense protoplasts (fig. 13,A, B). Each protophloem (or phloem) pole (six in fig. 13,A) has two or more sieve elements disposed tangentially (fig. 13,A, B).

Fig. 10. Electron micrographs of cortical root tip cells. Potassium permanganate fixation. A. Intercellular spaces (i) are almost filled with electron-dense material. B. Tannin vacuoles. At arrow, tangential section of a tannin vacuole. Details: arrow, tannin; d, dictyosome; er, endoplasmic reticulum; i, intercellular space; m, mitochondrion; n, nucleus; pl, plastid; v, vacuole; w, cell wall.

Fig. 11. Electron micrographs of cortical root tip cells. Potassium permanganate fixation. A. Cell with large vacuole containing tannin. Parietal cytoplasm with numerous endoplasmic membranes is sharply delimited because of tannin accumulation at periphery of vacuole. B. Crystal cell with coiled agglomeration of endoplasmic reticulum in parietal cytoplasm. The electron-transmitting sites of raphide crystals (r) are confined to vacuole (v). Black angular bodies, probably sections of crystals themselves. Details: d, dictyosome; er, endoplasmic reticulum; i, intercellular space; m, mitochondrion; r, site of raphide crystal; v, vacuole; w, cell wall.

Fig. 12. Electron micrographs of procambial cells from longitudinal sections of root tip. Potassium permanganate fixation. A. Lobed nucleus at n. B. At m extensive mitochondrion with branched profile—probably a partial section of one having the shape of an umbo (see also fig. 8). Details: er, endoplasmic reticulum; m, mitochondrion (both letters are in the same mitochondrion); n, nucleus; w, cell wall.

Fig. 13. Root tip sections (transverse in A, B, longitudinal in C) from paraffin-embedded material. A. Central cylinder and part of cortex at level where the first sieve tubes are mature. There are six phloem poles. B. Enlarged view of one of the phloem poles with sieve tubes at s. C. Part of cortex with longitudinal series of raphide crystal cells in different stages of development. First indication of development of such cell is indicated by check mark. Details: check mark, youngest identifiable raphide cell; en, endodermis; r, raphide cell; s, sieve element. (A, X150; B, X1,000; C, X125.)

Fig. 14. Differentiating sieve tube from longitudinal section of root tip. Paraffin embedding. Seventeen sieve tube members, some mature but mostly immature, are shown, with the youngest at 1. Cell 10 is distinctly elongated and contains a slime body above the nucleus. Nuclei are disintegrating in cells 13 and 14. Curved body in cell 14 is a slime body. Cells 15 to 17 are enucleate and their slime bodies are disaggregated. (X1,300.)

The vertical files of protophloem sieve elements observed in longitudinal sections reveal the successive stages of differentiation of the elements (fig. 14). The sieve elements are at first quite short and have dense protoplasts; elongation and increase in vacuolation follow. A slime body may be discerned at this time (cell 10 in fig. 14). Just before elongation occurs the protoplast seems to become especially sensitive to injury, as is indicated by its frequently unsatisfactory fixation (cells below cell 10 in fig. 14). Elongation and vacuolation progress, nuclei disintegrate, and slime bodies disperse. The sieve element assumes the characteristic clear aspect (cell 15 and those above it in fig. 14).

The clearness of sieve element protoplasts is also evident in electron microscope preparations (s in fig. 15,A, and fig. 16,B). The cells contain a thin parietal layer of cytoplasm with some organelles. No tonoplast is discernible. According to other studies on phloem (Esau and Cheadle, 1965), the space formerly occupied by the vacuole contains mictoplasm (Engleman, 1965), a derivative of the vacuole and the protoplasmic material that is released after breakdown of the tonoplast. With the fixations used in the present study, the mictoplasm contains some electron-dense material in the form of a thin irregular network or as discontinuous flocculent aggregations.

The cells associated with the sieve elements have rich complements of organelles and membrane systems and a relatively dense hyaloplasm (fig. 15,A). Some of these cells contain tannins, others have tannin-free vacuoles, and their nuclei may be lobed (figs. 15,A, and 17,C, at n in both). The dictyosomes are associated with numerous vesicles. The accumulations of endoplasmic membranes are particularly remarkable (fig. 17,A) and these occur not only in elongated cells of the vascular region but also in shorter, probably pericyclic cells (fig. 17,C). Thus,

cells with such accumulations may or may not be differentiating sieve elements. The youngest sieve elements recognized are exemplified by the cell in median position in figure 16,A (see also left side of fig. 15,B). The sieve element depicted has a nucleus, plastids, mitochondria, and few dictyosomes. The endoplasmic reticulum is sparse and partly broken down into vesicles (ve). The nuclear envelope also shows indications of a separation into vesicles. In later stages, vesicles are the dominant form of endoplasmic reticulum (fig. 16,B), although sometimes concentric aggregations of double membranes are encountered in enucleate cells (figs. 17,B, and 24,E). These aggregations are distinct from the slime, which is fibrous in appearance (fig. 21,C). Plastids and mitochondria are maintained (fig. 16,B). The material proved inadequate for study of developmental changes of these organelles. Available sections indicated no obvious differences between

the mitochondria of sieve elements and those of the contiguous cells.

As mentioned in "Materials and Methods," all sieve areas observed contained electron-opaque material in their pores, probably largely slime. In view of the controversy regarding interpretation of pore contents in sieve areas (Esau and Cheadle, 1965), the dense material in the pores is here referred to as protoplasmic pore contents or protoplasmic strands. Slime is of protoplasmic origin, and therefore the word "protoplasmic" does not exclude slime.

The strands in the sieve plates of protophloem are somewhat thicker than plasmodesmata (figs. 17,B, and 18,B) and are associated with callose (fig. 18,A). They are quite distinct in cells that still have nuclei (one is visible in cell s in fig. 15,B). At the end of the functioning period of the sieve element definitive callose is deposited. It may show fine, presumably protoplasmic strands (fig. 18,C).

SECONDARY PHLOEM IN THE STEM

Contents of sieve elements

The protoplast of *Vitis* sieve element, as seen with the light microscope, has been studied in considerable detail previously (Currier, Esau, and Cheadle, 1955; Esau, 1948a). In view of the sustained controversy regarding the nature of the sieve element protoplast in general and of one of its components, slime, in particular (see "Discussion") the behavior of this substance in *Vitis* was re-examined.

In differentiating sieve elements of *Vitis* the slime forms discrete bodies variable in shape and size. In constitution the bodies are frequently amorphous in killed material (fig. 19,G) but may show a regular pattern in arrangement of component elements (fig. 19,H). The dispersal of the bodies coincides with nuclear breakdown (fig. 19,I). After the slime bodies are disintegrated

the protoplast assumes various forms in killed material. Commonly, the sieve plates bear accumulations of slime (slime plug) that fill the pores (fig. 20,E; see also Esau, 1948a). Within the lumen of the cell the protoplast is contracted but is firmly attached to one or both sieve plates. Such attachment is particularly conspicuous in the short sieve elements that traverse the rays (fig. 20,A). The inner part of the protoplast may appear stringy, with more or less distinct strands merging with the pore contents of the sieve plates (fig. 20,B-D), or it may compose one continuous mass (fig. 20,E).

Because the protoplast is contracted in all these views they cannot be relied on for the interpretation of the normal uninjured condition of the sieve element contents. In studies on plasmolysis in *Vitis* phloem (Currier, Esau, and Cheadle, 1955) the presence of strands,

Fig. 15. Electron micrographs of phloem region in root tips in transverse (A) and longitudinal (B) sections. Potassium permanganate fixation. A. Phloem pole from second mm from apex. Cell between two mature sieve elements (s) contains tannin vacuoles. Cell with lobed nucleus (n) has a rich complement of endoplasmic reticulum. Cells with large vacuoles at v. B. Phloem pole from third mm from apex. Mature sieve tube with sieve plate at pt. The larger organelles are plastids, the smaller are mitochondria. Immature sieve element (s) with nucleus (n) to the left. Details: n, nucleus; pt, sieve plate; s, sieve element; tv, tannin vacuole; v, vacuole; w, cell wall.

Fig. 16. Electron micrographs of phloem from longitudinal sections of root tips. Potassium permanganate fixation. A. Immature sieve element from third mm from apex (see fig. 15, B). Nuclear envelope is discontinuous—possibly in early stage of disintegration into vesicles. B. Enucleate sieve element from second mm from apex. Endoplasmic reticulum and probably the nuclear envelope also have disintegrated into vesicles. Details: d, dictyosome; m, mitochondrion; n, nucleus; pl, plastid; pt, sieve plate; ve, vesicles; w, cell wall.

Fig. 17. Electron micrographs of phloem region from root tip in longitudinal sections. Potassium permanganate fixation. A. Part of immature sieve element from third mm from apex and adjacent procambial cell with dense accumulations of endoplasmic membranes. These membranes are partly vesiculated in the sieve element. B. Parts of mature sieve elements with sieve plate (pt) between them from first mm from apex. The coiled aggregation consists of double membranes (see also fig. 24, E). C. Cell from phloem pole, possibly pericyclic, from third mm from apex. Conspicuous accumulation of endoplasmic membranes and a lobed nucleus. Details: d, dictyosome; er, endoplasmic reticulum; er?, probably endoplasmic reticulum; n, nucleus; pl, plastid; pt, sieve plate; w, cell wall.

Fig. 18. Electron micrographs of sieve plates from root tip in transverse (A) and longitudinal (B, C) sections. Potassium permanganate fixation. A. Part of sieve plate (pt) in face view. Electron-dense material in pores surrounded by callose (white). B. Sectional view of plate. Callose (ca) covers plate on both surfaces and encloses strands. C. Sieve plate embedded in definitive callose traversed by thin strands of probably protoplasmic material. Sieve plate itself is not discernible. Details: ca, callose; pt, sieve plate; w, cell wall.

Fig. 19. Cytokinesis and protoplasts of sieve elements and companion cells in longitudinal sections of secondary phloem. Chrome-acetic-formalin fixation and mounting in Karo syrup. Collected May 30, 1945; photographed October 30, 1964. A. Phragmoplast and cell plate in parenchyma cell. B. Part of sieve-element mother cell in process of forming a companion cell. Nucleus (cut on bias) in mitosis. Phragmosome outlines the future companion cell. C, D. Com-

Fig. 20. Sieve element protoplasts in longitudinal sections of secondary phloem. Chrome-acetic-formalin fixation and mounting in Karo syrup. Collected, A, E, May 30, 1945; B-D, September 18, 1946. All photographed October 30, 1964. A. Commissural sieve elements crossing a ray. Their protoplasts are stringy (st) and continuous from cell to cell across the sieve plate (pt). B-D. Stringy contracted protoplasts attached to sieve plates. E. Slime plug showing differentiation between parietal cytoplasm and enclosed slime—part of the microplasm. Dense slime in sieve plate pores. (All, X840). Details: ct, cytoplasm; pt, sieve plate; sl, slime; st, stringy protoplast.

panion cells with slime bodies (sl). E. Companion cell with almost entirely dispersed slime body (sl). F. Mature companion cell with plastid-like organelles (or). G, H. Sieve elements with two forms of slime bodies, one dense (G), the other loose and fibrous (H). I. Sieve element with disorganizing nucleus (n) and disaggregating slime (sl). (A-G, X1,400; H, X840; I, X560.) Details: cp, cell plate; mt, nucleus in mitosis; n, nucleus; or, unidentified organelles; ph, phragmoplast; phs, phragmosome; sl, slime or slime body; w, wall between companion cell and sieve element.

Fig. 21. Electron micrographs from secondary phloem (A, B) and root tip (C). A. Tangential longitudinal section including sieve element (s) and companion cell (cc) from dormant cane collected December 12, 1960. Potassium permanganate fixation. Slime aggregation (sl) is fibrous in appearance. Starch grains at g. The wall between companion cell and sieve element with four pit-fields penetrated by plasmodesmata (see only in part). Numerous mitochondrion and nucleus in companion cell. B. Slime in sieve element from dormant cane. Osmium fixation. C. Slime in sieve element from root tip after fixation with potassium permanganate. Details: arrows, possibly fat bodies; cc, companion cell; g, starch grain; m, mitochondrion; n, nucleus; s, sieve element; sl, slime; 1, 2, 3, 4 are primary pit-fields.

including longitudinally-oriented ones, was found to be an indication of change leading to loss of plasmolyzability.

Among the several components encountered in the sieve elements of *Vitis* by means of the electron microscope, only one could be correlated with views of slime under the light microscope. As shown in the earlier study (Esau, 1948a), dormant sieve elements frequently contain dense aggregations of slime that may be continuous through occasional pores left unoccluded by dormancy callose (plate 20, D, in Esau, 1948a). Dense aggregations of slime were observed in dormant material collected for electron microscopy. Permanganate fixation indicated a faintly fibrous condition of this slime (fig. 21,A). Osmic acid treatment made the slime appear coarsely fibrous (fig. 21,B). Continuity of slime through pores in dormant cells was again observed. The strands appeared more or less fibrous in sections killed in osmic acid (fig. 22,C and E). Some dormant sieve elements revealed a fine net of fibrils after treatment with formalin and KMnO_4 (fig. 22,D).

Two formations encountered in *Vitis* sieve elements under the electron microscope have not been properly identified. One of these, found before and during dormancy, consisted of dense aggregations of parallel-oriented membranes (fig. 22, A and B). Small circular profiles occurring near the margins of these aggregations (fig. 22,A) suggested that the membranes may pertain to the endoplasmic reticulum, tubular in form. Mehta and Spanner (1962) depicted similar material in sieve elements of *Nymphoides peltatum* and interpreted it as endoplasmic reticulum.

The second unidentified element was seen in newly differentiated sieve elements. It appeared to be composed of single-layered membranes concentrically coiled in a manner suggesting the so-called myelin figures (fig. 25,C). The origin of this formation is obscure, but

its resemblance to the suspected coiled endoplasmic membranes in sieve elements of the root tip (fig. 24,E) suggests the possibility that it is derived from such membranes.

Sieve elements of *Vitis* contain plastids with starch (fig. 23,A) showing the familiar red coloration when treated with iodine. One or more granules occur in one plastid and they may occupy most or only part of a plastid (figs. 23 and 35,B). The plastids have double outer membranes and small, sparse inner membranes not assembled into grana (fig. 23,A–C). The starch grains are often only slightly electron opaque (fig. 23, A and C). When they appear darker, a concentric layering of the starch is revealed (fig. 35,B). No starch was encountered in the plastids in root tip phloem (fig. 24,E).

Companion cells

In the cytokinesis leading to the delimitation of a companion cell, a phragmosome in the sense of Sinnott and Bloch (1941)—that is, a cytoplasmic diaphragm—precedes the division of nucleus and the formation of cell plate (fig. 19,B). The phragmosome is clearly distinguishable from a cell plate in that its profile is not a sharp and rather straight line, as is that of the cell plate (fig. 19,A). The phragmosome indicates the shape of the future companion cell, elongated and tapering at the ends (compare fig. 19,B, with fig. 3).

As reported previously (Esau, 1947, 1948a), a companion cell in *Vitis* develops a single slime body resembling those in sieve elements (fig. 19,C and D). The slime body later disperses (fig. 19,E). As the tonoplast of the companion cell does not break down, the slime apparently remains incorporated in the cytoplasm, which assumes a denser appearance than it had before the slime body disintegrated.

Small plastid-like bodies become evident in the maturing protoplast (fig. 19,E and F at **or**). The bodies are num-

Fig. 22. Electron micrographs of details of sieve element protoplasts from longitudinal sections of canes. A. Collected October 20, 1961. Potassium permanganate fixation. Grains of starch at g and possibly endoplasmic membranes at er?. B. Collected February 9, 1962, induced reactivation during 2 weeks in greenhouse, and sampled February 23, when cane developed shoots less than 1 inch long. Potassium permanganate fixation. Starch grain at g and possibly tubular endoplasmic membranes at er?. C. Pore with slime during dormancy. Collected December 20, 1960. Osmic acid fixation. D. Mictoplasm of dormant sieve element. Collected January 25, 1962. Formalin and potassium permanganate fixation. E. Slime in pore during dormancy. Collected December 20, 1960. Fixation as in C, except that it was done on ice. Details: ca, callose; g, starch grain; er?, probably endoplasmic reticulum; w, cell wall.

Fig. 23. Electron micrographs of plastids in sieve elements. Fixations: A, C, D, potassium permanganate; B, osmic acid. Plastids contain one or more starch grains (g). Inner membranes are sparse and vesicular if present (A, B). Evidence of doubleness of outer membrane in B. Concentric layering in starch grains is faintly indicated in A and B. Details: arrow, double outer membrane; g, starch grain; er, endoplasmic reticulum.

erous and discrete in newly differentiated companion cells (fig. 19,F) but are replaced by fewer and larger entities resembling "deeply staining flakes" (Esau, 1948a, p. 244) in dormant state. Osmic acid fixation reveals black-stained globules comparable to the plastid-like bodies in size (fig. 24,B). It is not certain, however, whether the two kinds of bodies are identical.

Previously the suggestion was made that the plastid-like bodies are "left" after the slime is dispersed (Esau, 1948a, p. 241)—an idea that needs further examination. Companion cells with slime bodies were not encountered in electron-microscope material, and the organelles comparable in shape and size to the plastid-like bodies seen in that material had a structure not previously described for inclusions in plant cells. As seen after permanganate fixation, the bodies contained concentric lamellations surrounded by a membrane associated with vesicles in orderly arrangement (figs. 24,D, and 25,A and B). Some were intricately partitioned. Fixation was not sufficiently critical to determine whether any of the membranes were double. The lamellated organelles are considerably larger than mitochondria (fig. 25,B); they possibly are plastids modified by some growth conditions or by some part of the processing during preparation for electron microscopy. It seems doubtful that these organelles are related to the concentric aggregations of apparently endoplasmic membranes (fig. 24,E) or the "myelinated" structures (fig. 25,C) observed in sieve elements.

The organelles in the companion cells resemble the lamellated inclusions in fibroblasts of otocysts from fowl embryos cultured *in vitro* and exposed to the effect of Dimycin (Friedman and Bird, 1960). The authors found evidence that the inclusions were mitochondria changed degeneratively under the influence of Dimycin. One might also compare the organelles in the com-

panion cells with the myelinated fibers of the sciatic nerve of frog depicted by Robertson (1960).

The lamellated organelles of companion cells were found in newly differentiated phloem in a young shoot that developed on a cane reactivated in a greenhouse (see "Materials and Methods"). In dormant phloem treated with KMnO_4 numerous mitochondria and some small bodies with dark contours and light centers were present (fig. 21,A, unlabeled arrows), but no plastids were observed. The darkly contoured bodies could conceivably be the so-called fat bodies (Manton, 1962), which are incompletely fixed in KMnO_4 . If this suggestion is correct, the black globules in figure 24,B, may be identical with these fat bodies although there appears to be a discrepancy in size.

Some companion cells had moderate amounts of endoplasmic reticulum (fig. 21,A); others contained dense accumulations (fig. 26, A–C) resembling those previously described for cells associated with sieve elements in the root tip. These membranes appeared mostly as ribbons (fig. 26,C), but they were interpreted as endoplasmic membranes because in some views double-layered profiles were recognized (fig. 26,B). They appear to be tubular membranes. In one collection large accumulations resembling myelin figures were present (fig. 26,D). According to Friedman and Bird (1960), in the *in vitro* cultures of fowl embryo otocyst not only mitochondria degenerated but endoplasmic membranes also underwent swelling and "morphological alterations." The material in figure 26, D, could have resulted from modifications, including swelling, of aggregations of tubular endoplasmic membranes. Menke (1959) observed myelin figures in occasional shoot apices of *Elodea*. He was unable to identify these figures with any specific organelles or membrane systems but suggested that they indicate some degenerative change, perhaps related to growth inhibition.

Fig. 24. Photographs from monitor slides (A, B) and electron micrographs (C-E). A-D, companion cells from longitudinal sections of secondary phloem; E, sieve element from root tip. Fixations: A, chrome-acetic-formalin followed by staining with OsO_4 ; B, osmium; C-E, potassium permanganate. A. Deeply pitted wall (w) is between companion cell and sieve element. (X1,400). B. Osmiophilic granules and pitted walls. Sieve element was to the right. (X1,800.) C. Pit areas are provided with plasmodesmata connecting sieve element and companion cell protoplasts. D. Cell from young phloem. Coiled material ("myelin figures") apparently within plastids. (See also fig. 25.) E. Plastids and double coiled membranes, possibly endoplasmic. (See also fig. 17, B.) Details: er?, probably endoplasmic reticulum; m, mitochondrion; n, nucleus; pi, pit-field with plasmodesmata.

Fig. 25. Electron micrographs of companion cells (A, B) and sieve element (C) from longitudinal sections of young secondary phloem. Potassium permanganate fixation. A, B. Coiled material and vesicles possibly within plastids. (See also fig. 24, D.) Intact mitochondria in B. C. Coiled material and starch grains (g) in sieve element. Details: g, starch grain; m, mitochondrion; w, cell wall.

Fig. 26. Electron micrographs of contents in companion cells from longitudinal sections of secondary phloem. Potassium permanganate fixation. A–C. Aggregations possibly composed of endoplasmic membranes. They appear double at arrows in B. At check marks in B, possibly sections of membranes. D. Material in companion cell from dormant cane resembles that in companion cells in figures 24 and 25. Details: arrows, double membranes; check marks, circular profiles; er?, probably endoplasmic reticulum; w, cell wall.

Fig. 27. Electron micrographs of sieve areas seen in surface views, from longitudinal sections of secondary phloem collected October 20, 1961. Potassium permanganate fixation. A. Part of sieve area from sieve plate in end wall, as seen in radial section. Pores filled with electron-dense material surrounded by callose (ca). B. Sieve areas in radial longitudinal wall. Pore contents and callose as in A but of smaller dimensions. Details: ca, callose; w, cellulose part of cell wall.

Authors are commonly reluctant to suggest fixation artifacts. According to Ito (1962), myelin figures may arise as a result of post-mortem changes of cytoplasmic membranes (example, plasmalemma in bat kidney tubule), but in general cellular membranes are more resistant to such changes than is commonly supposed. He thinks that artifacts of specimen fixation are sometimes interpreted as indications of post-mortem autolysis. Some of the myelinated formations in *Vitis* may be such artifacts but, conceivably, information about the factors inducing their formation may increase our understanding of the normal structures.

The wall between the companion cell and the sieve element is deeply pitted (primary pit-fields, figs. 19,F, and 24,A and B). The pit membranes are traversed by plasmodesmata (figs. 21,A, and 24,C). Indication of branching of plasmodesmata on the companion cell side (Esau and Cheadle, 1965) was noted in *Vitis* phloem.

Sieve plates and lateral sieve areas

The characteristic inclination and compounding of sieve plates of *Vitis* are seen in figure 3 at **pt**. In size of pores the sieve areas of the sieve plates (fig. 27,A) intergrade with sieve areas on the side walls (fig. 27,B). In all material processed for this study the pores in active phloem were filled with electron-dense material, probably slime. Furthermore this material projected, in the form of radially-oriented fine strands, into the callose lining the pores (fig. 27).

Views of developmental stages of sieve areas reveal features similar to those described for *Cucurbita* (Esau and Cheadle, 1965). Each pore site bears two callose platelets opposing one another on the two sides of the wall (fig. 28,B). Surface views of sieve plates in this stage of development show that each pore site has one plasmodesma (fig. 28,A). This observation agrees with the

previous interpretation, made by use of the light microscope, that in *Vitis* the number of plasmodesmata in the primary pit-fields of cambial walls corresponds to the number of connecting strands in the sieve areas of a sieve plate (Esau, 1948a, p. 234). Partial longitudinal sections of plasmodesmata in young sieve areas are discernible in figures 28, B, D, and E. The dark cytoplasmic material on the pore sites in figure 27 does not reveal the relation of endoplasmic reticulum to pore sites observed in *Cucurbita* (Esau and Cheadle, 1965).

Dormancy callose occurs on sieve areas of various degrees of differentiation (fig. 29) but is particularly massive on the sieve plates (figs. 29,A, and 30,A). It usually covers not only the individual sieve areas but also the thickened bars of wall intervening between contiguous sieve areas (fig. 29,A and C). In sieve areas connecting the sieve elements with parenchyma cells, dormancy callose was observed only on the side of the sieve element (fig. 31,A; compare with plate 1, C and D, in Esau, 1948a).

Light-microscope studies gave uncertain results regarding presence or absence of protoplasmic strands in dormancy callose (Esau, 1948a, p. 243). Use of techniques for demonstrating plasmodesmata suggested that thin strands may be present, and electron microscope views support this assumption (fig. 30,A). The strands in the lateral sieve areas may be associated with median nodules (fig. 29, C and D).

The removal of callose during reactivation of sieve elements as seen with the light microscope suggests a dissolution process in which slime is involved. First, canals filled with slime appear in the callose (Esau, 1948a, plate 20,A), then this carbohydrate is removed from the entire surface of the plate. Ultrastructural views give a similar impression. Callose pads in the process of being removed show cavities filled with elec-

Fig. 28. Electron micrographs of immature sieve areas with closed pore sites from longitudinal sections of secondary phloem collected July 18, 1961. Potassium permanganate fixation. A Radial section exposing callose platelets (ca) at pore sites (arrows). One plasmodesma in each pore site (arrows). B. Oblique radial section exposing platelets in sectional view. Part of a plasmodesma at arrow. C-E. Tangential sections showing distribution of pore sites (numbered) with paired callose platelets. Parts of plasmodesmata at arrows. Details: arrows, plasmodesmata; ca, callose; numbers, pore sites.

Fig. 29. Electron micrographs (A, C, D) and photograph from a monitor slide (B) illustrating massive callose (ca) seen in sections of sieve areas. Potassium permanganate fixation. A. Dormancy callose occluding pores (numbered 1-4) and enclosing thinner and thicker wall parts of the sieve area of a sieve plate. Pores 2-4 constitute one sieve area. B. Dormancy callose is layered. (X1,400.) C. Sieve areas with dormancy callose from a side wall (not a sieve plate; a lateral sieve area). Protoplasmic material in the form of a strand (check mark) traversing one pore. D. Lateral sieve area with considerable amount of protoplasmic material—strands and median nodules—present in the callose. Details: ca, callose; g, granule of starch enclosed in plastid; po, pore in sieve area; w, wall of sieve plate, cellulosic part; check mark, protoplasmic strand; numbers, pores in sieve area.

Fig. 30. Electron micrographs of parts of sieve plates from tangential sections of secondary phloem. Potassium permanganate fixation. A. Dormancy callose with fine protoplasmic strands from a cane collected November 28, 1960. B. Dormancy callose in process of dissolution. From a cane collected February 9, 1960, placed in a greenhouse for 2 weeks to induce reactivation, and fixed February 23, 1960. Electron-dense protoplasmic material in pores within callose is largely slime. Arrow points to lighter-appearing callose, probably affected by enzyme action. Details: ca, callose; po, pore; w, wall of sieve plate, cellulosic part.

tron-dense material, probably slime. Significantly, the callose in immediate contact with the slime (unlabeled arrow in fig. 30,B) appears to have a different consistency than the still unaffected callose. An enzyme action is indicated by such views.

The canals are formed in the callose pads in line with the pores in the cellulosic part of the wall. Whether the fine protoplasmic strands remaining in the callose during dormancy (fig. 30,A) play any role in initiating or directing the process of dissolution of callose is not known.

In addition to the sieve areas just described—structures that were similarly interpreted at both the light- and electron-microscope levels—sieve areas with distinctive ultrastructure were recognized. These are minor, or least differentiated, lateral sieve areas between contiguous sieve elements. One such sieve area is shown in figure 31,C. Its strands, surrounded by callose (**ca**), hardly surpass in width the plasmodesmata in the primary pit-fields of parenchyma cells as depicted in figure 31,B. As far as could be ascertained, the callose of the minor sieve areas is not continuous through the wall but appears as two conical masses tapering toward the middle lamella region (fig. 32,B). Moreover, a distinctive kind of median nodule was observed in the preparations used: a more or less spherical structure lined with electron-dense material (fig. 32,C). The two features just described are seen in figures 31,C, and 32,A, both of which depict sections passing obliquely through the wall and revealing the same sieve areas at different depths within the wall. In figure 31,C, the section passes part way through the wall and demonstrates the diminution of diameters of the callose cones in the downward direction; open median nodules (**mn**) are seen below. In figure 32,A, the callose cones diminish in size toward the median part of the figure where median nodules (**mn**) occur, and

below these are the callose cones of the opposite side of the wall, appearing narrower near the median nodules and wider in the downward direction.

The minor sieve areas show still another feature not present in the more highly differentiated areas. Some callose cones contain more than one strand (fig. 32,A and D). This configuration results when two strands occur close together. It is possible to recognize callose cones and their strands in various degrees of approximation to one another (1–5 in fig. 31,C). The strands may be completely apart, each with its pair of callose cones (fig. 32,E), or they may occur so close to one another that the callose embeds two strands jointly (fig. 32,D).

In the 1948 study of *Vitis* phloem the assumption was made that sieve areas of the sieve plates and those of the side walls differ only quantitatively and that the two kinds of structures intergrade (Esau, 1948a, p. 234). The sieve areas of the sieve plates were designated sieve areas *A*, those of the side walls sieve areas *B*. The present study indicates that minor lateral sieve areas with discontinuous callose and prominent median nodules are also present.

Parenchyma cells and fibers

Axial parenchyma cells occur as parenchyma strands. Both axial and ray parenchyma cells store starch or contain tannins (figs. 3, 33, and 34,B). Sometimes starch and tannins occur in the same cells, starch in plastids in the cytoplasm, tannin in the vacuole. Starch grains develop one to several in one plastid and are usually quite light in electron micrographs (fig. 35,A); sometimes they disappear during preparation of material for study and leave spaces in the sections (fig. 33, A and B). The starch grains of parenchyma cells are clearly differentiated from the concentrically lamellated starch grains in the sieve elements (fig. 35,B).

Axial and ray cells are pitted (fig. 33, A and B). In the ray cells a thickening

Fig. 31. Electron micrographs of lateral sieve areas (A, B) and a primary pit-field (B). Potassium permanganate fixation. A. Longitudinal section passing through a sieve element (s) and a parenchyma cell (pa). Wall (w) between the two cells has dormancy callose (ca) on the side of the sieve element. B. Primary pit-field with plasmodesmata in parenchyma cell seen in face view in a radial wall. C. Lateral sieve area seen in face view in a radial wall. Tissue not dormant. Collected July 18, 1961. Sieve area forms a lateral connection between two sieve elements. From the upper right to the lower left the section passed at increasingly greater depth through the wall and is at middle lamella level at left. Here, the protoplasmic material (black) occurs in the form of median nodule (see fig. 32). One or two strands are encased in one callose (ca) cylinder. Numbers 1-5 indicate pairs of strands with various degrees of confluence of their callose jackets. Details: arrows, protoplasmic strands within common callose mass; ca, callose; mn, median nodule; pa, parenchyma cell; s, sieve element; w, cellulosic part of cell wall; 1-5, pore sites of different degrees of approximation to one another.

occurs on the pit membrane (fig. 33,D, unlabeled arrow) and the small intercellular spaces show a centrally located structure (fig. 33,D, at i), which has not been identified. The walls of parenchyma cells are here interpreted as primary. In some preparations they show a degree of heterogeneity, and tanniniferous cells have a lighter-staining wall than do those free of tannin (fig. 33,C).

Many tannin-free parenchyma cells contain abundant endoplasmic reticulum (fig. 34,A) as well as mitochondria. Tannins appear as irregular somewhat ramified masses spread through

the vacuole (fig. 33,E). Tannin cells have a thin layer of parietal cytoplasm connected by plasmodesmata with the protoplasts of contiguous cells (fig. 34,B).

The fibers, frequently septate, store starch and have moderately thick secondary walls. According to light-microscope studies the pits were interpreted as bordered (Esau, 1948a, p. 230). An indication of a border is seen in the electron micrograph in figure 36,A. As the fibers are living cells, plasmodesmata occur in the thin pit membranes (fig. 36,B) of the deeply-recessed pits.

DISCUSSION

The phloem of the thirty-one species of *Vitis* (including *V. vinifera*) examined in this study shows uniformity of structure extending even to such features as types of crystals and their distribution, occlusion of nonconducting sieve elements by tylosoids, and the tendency of a given phloem increment to function in conduction for more than one season.

The deep-seated origin of the first periderm was found to be characteristic of thirty species. In the thirty-first, *V. rotundifolia*, the first periderm arose close to the surface and was still in the same position in several year-old branches; that is, such branches still had then living (but nonconducting) primary phloem tissue and most of the cortex. In other species these two tissue regions were not present in branches in the second year of growth (canes) because the phellogen arose in the primary phloem. Moreover, in older branches the first periderm was replaced by periderms of still deeper origin. This developmental pattern is responsible for the rough, shredding appearance of the bark in *Vitis* (ringbark) other than *V. rotundifolia*. The latter species has a smooth nonshredding bark (Bailey and Bailey, 1949) because only a relatively small amount of tissue (cork) is

sloughed off as the stem grows older.

Vitis continues to be the genus best investigated with regard to the phenomenon of reactivation of phloem (see reviews on longevity of sieve elements in Esau, 1948a, pp. 218–219, and Evert, 1963). The course of development of the massive dormancy callose and the removal of this callose in the spring has been studied in detail with the light microscope; some information about the phenomena is available at the ultrastructural level. Both approaches indicate that slime is involved in the removal of callose.

Because the nature and normal state of slime in the sieve element are subjects of much research and discussion, it will be useful to review the behavior of the substance during the reactivation of phloem and, incidentally, to recall the earliest observation on the phenomenon by Wilhelm (1880).

Dormancy callose does not readily reveal any protoplasmic strands when examined with the light microscope. Fine strands may be seen occasionally after treatment of sections with stains used for demonstrating plasmodesmata, but the slime, which can be differentially stained in *Vitis*, is usually completely detached from sieve plates that are covered with dormancy callose (Esau,

Fig. 32. Electron micrographs of lateral sieve areas from nondormant secondary phloem collected July 18, 1961. They illustrate lateral connections between sieve elements. Potassium permanganate fixation. A. Sieve areas seen in face view. From above downwards, section passed from one wall surface through the middle lamella region (median nodule, mn, region) to the other wall surface. B, C. Oblique sections through pore sites including one of a pair of conical blocks of callose. In B, protoplasmic strand within callose. D. Double pore site comparable to the one within two arrows in A. E. Pair of pore sites comparable to number 2 in figure 31, C. D and E probably not median sections of nodules. Details: ca, callose; mn, median nodule; arrows, protoplasmic strands.

Fig. 33. Photographs from monitor slides (A-D) and electron micrograph (E) illustrating ray and parenchyma cells from dormant secondary phloem. Potassium permanganate fixation. A. Tangential section including axial parenchyma (pa) above and ray cells below. Both types of cells contain starch. B-D. Enlarged views of ray cells, some containing starch (B) others tannin (C). Prominent simple pits (pi) with thickening on the pit membrane (unlabeled arrow in D). In the tanniniferous cell in C the secondary wall is rather thick and lightly stained. Unidentified dark structure in intercellular space (at i in D). E. Tannin in a vacuole of a parenchyma cell. (A, X780; B, C, X1,400; D, X1,800.) Details: unlabeled arrow, thickening on pit membrane; cr, crystals; g, starch granule; i, intercellular space; pa, parenchyma cell; pi, pit; ta, tannin; w, cell wall.

Fig. 34. Electron micrographs of axial parenchyma cells from dormant secondary phloem. Potassium permanganate fixation. A. Endoplasmic reticulum (er) is abundant and plasmodesmata are present in the wall (w). B. Lower cell has thin layer of parietal cytoplasm (cy) and a tannin-filled vacuole. Plasmodesmata connect this cell with one containing no tannin. Details: cy, parietal cytoplasm; er, endoplasmic reticulum; pi, pit; w, cell wall.

Fig. 35. Electron micrographs of cells from secondary phloem. Potassium permanganate fixation. A. Axial parenchyma cell with plastids containing one or more starch grains (g). B. Plastids with starch grains from a sieve element. Details: g, starch granule; n, nucleus; ta, tannin; w, cell wall.

Fig. 36. Electron micrographs of parts of fibers illustrating secondary walls with pits and plasmodesmata. A. Barium permanganate fixation. B. Potassium permanganate fixation; protoplast withdrawn from cell wall. Details: pi, pit; w2, secondary wall.

1948a). In exceptional sieve plates some pores are incompletely occluded and slime is clearly continuous from cell to cell through such pores (Esau, 1948a, pl. 20,D). Incidentally, *Tilia* regularly shows partly-open pores in dormant sieve elements (Evert and Derr, 1964a).

In dormancy callose of *Vitis*, electron microscopy discloses fine strands, mostly appearing discontinuous in sections (figs. 29, C, D, and 30, A), and also occasional thick strands presumably composed largely of slime (fig. 22, C and E). Hepton and Preston (1960) report having recognized continuous strands in the callose of dormant sieve elements of *Vitis*, but their samples were taken from rooted cuttings in which cambial activity was resumed. One may question whether Hepton and Preston (1960) were dealing with unaltered dormancy callose, because when cambium is active phloem reactivation is under way (Esau, 1948a).

Dormancy callose frequently shows layering in both light- and electron-microscope views (figs. 29,B, and 30,A). In some preparations for electron microscopy considerable damage to tissue occurred during methacrylate polymerization. In such material cellulose walls and callose both were separated into layers. Layering of dormancy callose of *Vitis* was recorded in my previous study (Esau, 1948a) and was recognized also by Kessler (1958) and Wilhelm (1880, figs. 80 and 85). In the California material of *Vitis*, dormancy callose was found to be thicker than the definitive callose (callose appearing before complete cessation of sieve element activity) and the latter showed no layering (Esau, 1948a). Because dormancy callose also may lack conspicuous layering, Hepton and Preston's (1960) statement that dormancy callose in *Vitis* shows no layering is probably based on too-limited observations. Coarse layering of dormancy callose was seen by Evert and Derr (1946a) in electron micrographs of *Tilia americana* sieve elements.

The cause of layering in callose is obscure. According to Eschrich (1956), callose has a completely homogeneous structure at the ultrastructural level and is anisotropic (contrary observation by Küster, 1951). Possibly, the layering is caused by other substances deposited with the callose. Paolillo (1963) suspects presence of protein deposits in the callose of *Isoetes* phloem, and Eschrich (1962) suggests that components of cytoplasm occur at least temporarily, in the callose of pollen mother cell walls in *Cucurbita ficifolia*.

During reactivation callose is removed gradually. At first, thin strands crossing the callose and passing through the pores in the sieve plate become discernible (Esau, 1948a; Wilhelm, 1880). Later the strands increase in thickness, first at the periphery of the callose pads, then also at greater depth. This phenomenon has been interpreted as a penetration of callose by slime (Hill, 1908; Esau, 1948a). Views of sieve plates at this time suggest a dissolution of callose as seen with the light microscope (Esau, 1948a, pl. 20, A) and in electron micrographs (fig. 30,B). The change in the ultrastructural appearance of callose next to the strands has been interpreted as indicating that an enzyme is involved in the process (Hepton and Preston, 1960). Eschrich (1961) has obtained callose-degrading enzyme preparations from phloem exudate of *Cucurbita*. Whether such enzyme originates in the slime needs to be investigated.

The present study furnishes a confirmation of the origin of slime in the form of discrete entities in the cytoplasm of sieve elements and companion cells of *Vitis*. As in *Cucurbita* (Esau and Cheadle, 1965) the slime bodies in *Vitis* become dissociated as such at the time of nuclear degeneration in the sieve elements. The two phenomena are not causally connected, because in the companion cell the slime body disperses in the presence of an unaltered nucleus. In the

sieve element, the disappearance of the tonoplast results in an interpenetration of cytoplasmic and vacuolar contents so that the slime comes to lie in the interior of the cell rather than in the peripheral cytoplasm. The mixture is the mictoplasm as defined by Engleman (1965).

The conformation of slime in the enucleate, tonoplast-free cell is variously interpreted. Eschrich (1963*a,b*) depicts the mictoplasm as a net which he calls lipo-proteid net. Esau and Cheadle (1965) draw attention to the variable fixation images of the mictoplasm and slime in *Cucurbita*. The slime frequently has a fibrous appearance (Engleman, 1963).

Sieve elements may contain a fine or coarse network or straight strands of protoplasmic material extending from plate to plate. Both formations result presumably from presence of slime in the mictoplasm. The stringy condition of sieve element protoplasts is a phenomenon familiar to phloem investigators and has been recognized in the earliest studies on the tissue (Wilhelm, 1880, figs. 96 and 106). In our plasmolytic studies (Currier, Esau, and Cheadle, 1955) the cells with longitudinally-oriented strands and stringy networks were identified as injured cells. Kollmann (1960*a*) also interprets the strands as artifacts resulting from injury to the cytoplasm. He points out the important fact that not infrequently the strands become visible only after a treatment with a fixative (a mild fixative, dilute potassium iodide, was used). However, Evert and Derr (1964*b*) think that the strands are normal constituents of sieve element protoplasts, and Evert and Murmanis (1965) insist that the fibrillar material, which they recognized with the electron microscope, represents components of the strands seen with the light microscope. Evert and Derr (1964*b*) also saw strands in fresh material but less frequently than in the killed. Their contention is that the strands easily break down but are

preserved by rapid fixation. Thaine (1964) also advances the idea that the strands in sieve elements are normal structures. He shows these strands in severely contracted protoplasts and in the phloem exudate of *Cucurbita*, and interprets them as transcellular strands of cytoplasm. Although the Evert and Derr (1964*b*) study was carried out with due appreciation of the difficulties involved in the preservation of sieve element protoplasts, it has not ruled out the probability that the strands appear in response to treatment with fixatives or because of some other injury. The formation of strands may be as sensitive a reaction as the formation of callose (Evert and Derr, 1964*a*). If, however, the phenomenon was a response to injury, the resulting sudden appearance of strands extending from one end of the cell to the other would need an explanation.

The literature on the ultrastructure of the sieve element protoplast is extensive but it does not allow as yet the drawing of a complete picture of this aspect. We have reviewed the pertinent articles and pointed out areas of agreement and disagreement among investigators (Esau and Cheadle, 1965). One of the controversial matters is the condition of mitochondria. In *Cucurbita* these organelles show a depauperation of internal membranes before the sieve plate is perforated. In other plants they appear to retain their original structure in mature sieve elements (*cf.* Kollmann and Schumacher, 1964). In *Vitis* the material used was not suitable for drawing definite conclusions.

The other protoplast component receiving divergent interpretations is the endoplasmic reticulum. We have previously emphasized that in permanganate-fixed material this membrane system appears dissociated into vesicles when the sieve element is well differentiated (Esau and Cheadle, 1965). Kollmann and Schumacher (1964) suggest that the vesiculation is induced by fixation,

although the response as such may be taken as an indication of an altered physiologic state of the cell. Vesiculation of endoplasmic membranes was observed in sieve elements of *Vitis* root tips. As a result these elements sharply contrasted with the surrounding nucleate cells, many of which had particularly strong development of endoplasmic reticulum. Some mature sieve elements had concentric aggregations of double membranes resembling those found in raphide crystal cells.

No developmental stages of sieve elements were obtained in the secondary phloem, but in some mature elements dense accumulations of linear entities resembling endoplasmic membranes were found. Similar accumulations occurred in companion cells. One might compare these accumulations with the massive aggregations of endoplasmic membranes depicted by Kollmann and Schumacher (1964) for *Metasequoia* sieve cells which occurred in a tissue killed *in situ*. The authors suggest that these accumulations might be the slime bodies observed by other authors. Tubular structure was not recognized in the

material identified as slime in *Cucurbita* (Esau and Cheadle, 1965) and *Vitis*, but we used different methods of fixation than did Kollmann and Schumacher.

The sieve element plastids of *Vitis*, which belong to the type that forms starch staining red with iodine, are remarkably poor in internal membranes. Similar observations were made by Kollmann (1960b) on *Passiflora* and by Kollmann and Schumacher (1964) on *Metasequoia*. The starch-forming plastids in phloem parenchyma cells of *Vitis* have more or less abundant grana (Esau, 1963).

Some features observed in this study with the electron microscope were not completely interpreted (for example, formations resembling myelin bodies in companion cells and sieve elements) but ultrastructural research is still at a stage in which natural phenomena and responses to fixations or other injuries are not always reliably distinguishable. Nevertheless, repetitive appearances of certain fixation images enable one to discuss cytoplasmic differentiations in terms of cell specializations.

SUMMARY

Thirty species of grapevine, *Vitis*, were compared for their bark characteristics and seasonal behavior of the phloem, the food-conducting tissue. This tissue was previously studied in detail in *V. vinifera*. In the present article that study is complemented by information obtained with an electron microscope.

The first periderm arises within the primary phloem in all species of *Vitis* except *V. rotundifolia*, in which it originates superficially. *V. rotundifolia* thus lacks the ringbark typical of the genus *Vitis*.

The phloem of *Vitis* differs from that in the majority of dicotyledons in its dormancy and reactivation. During dormancy the sieve plates are occluded by massive deposits of callose which, ultra-

structurally, are seen to enclose extremely fine protoplasmic strands. During reactivation, dormancy callose is removed by a process that, judged by electron microscope views, probably involves enzyme action. The proteinaceous material called slime possibly is involved in the removal of callose. Dormancy and reactivation phenomena were observed in almost all species of *Vitis* used in this study.

During the differentiation of sieve elements the pores of the sieve plates develop at sites penetrated by plasmodsmata, one to a pore site. Callose platelets cover the pore sites on both sides of the wall before the site is perforated. Sieve areas of various degrees of differentiation occur on the side walls. The

least differentiated ones have conspicuous enlargements of strands in the middle lamella region—the median nodules—and callose associated with these strands does not appear to be continuous across the wall.

Ultrastructural studies support the concept that the sieve element has a specialized protoplast. In addition to the well-known lack of nucleus and tonoplast (vacuolar membrane), it has an

endoplasmic reticulum which in fixed material appears dissociated into vesicles instead of forming extended cisternae such as are found in less specialized nucleate cells. The plastids of sieve elements elaborate starch that shows fine concentric lamellation at the ultrastructural level. The plastids have much less differentiated internal structure than do the plastids of parenchyma cells of the same tissue.

ACKNOWLEDGMENTS

The writer owes thanks to Dr. Klayton E. Nelson for having assisted with the preparation of slides for the study of *Vitis* species; to the Department of Viticulture, University of California, Davis, for financially supporting this aspect of research; to Mr. Edward B. Risley for technical assistance; and to Mr. Jack Pangborn for the introductory training in electron microscopy.

I appreciate the courtesy of the authorities of the California Agricultural Experiment Station in giving me the privilege of publishing in *Hilgardia* after my official connection with the Station was discontinued.

The study was supported in part by the National Science Foundation Grants G-14424 and GB-1523, and Public Health Service Grant R. G. 5868.

LITERATURE CITED

- BAILEY, L. H., and E. Z. BAILEY.
1949. Hortus second. New York: Macmillan.
- BELL, P. R., and K. MÜHLETHALER.
1964. The degeneration and reappearance of mitochondria in the egg cells of a plant. Jour. Cell Biol. 20:235-48.
- CHEAVIN, W. H. S.
1938. The crystals and cystoliths found in plant cells.
Part I: Crystals. Microscope, Brit. Jour. Micros. and Photomicrogr. 2:155-58.
- Committee on Nomenclature, International Association of Wood Anatomists.
1957. International glossary of terms used in wood anatomy. Trop. Woods 1957 (107):1-36.
- CURRIER, H. B., K. ESAU, and V. I. CHEADLE.
1955. Plasmolytic studies on phloem. Amer. Jour. Bot. 42:68-81.
- ENGLEMAN, E. M.
1963. Fine structure of the proteinaceous substance in sieve tubes. Planta 59:420-26.
1965. Sieve element of *Impatiens sultanii*. 2. Developmental aspects. Ann. Bot. 29:103-118.
- ESAU, K.
1944. Anatomical and cytological studies of beet mosaic. Jour. Agr. Res. 69:95-117.
1947. A study of some sieve-tube inclusions. Amer. Jour. Bot. 34:224-33.
1948a. Phloem structure in the grapevine, and its seasonal changes. Hilgardia 18:217-96.
1948b. Anatomic effects of the viruses of Pierce's disease and phony peach. Hilgardia 18:423-82.
1963. Ultrastructure of differentiated cells in higher plants. Amer. Jour. Bot. 50:495-506.
- ESAU, K., and V. I. CHEADLE.
1965. Cytologic studies on phloem. Univ. Calif., Publs., Bot. 36:253-344.
- ESAU, K., V. I. CHEADLE, and E. B. RISLEY.
1962. Development of sieve-plate pores. Bot. Gaz. 123:233-43.
- ESCHRICHT, W.
1956. Kallose. Protoplasma 47:487-530.
1961. Untersuchungen über den Ab- und Aufbau der Callose. (III. Mitteilung über Callose). Ztschr. f. Bot. 49:153-218.
1962. Elektronenmikroskopische Untersuchungen an Pollenmutterzellen-Callose. IV. Mitteilung über Callose. Protoplasma 55:419-22.
1963a. Beziehungen zwischen dem Auftreten von Callose and der Feinstruktur des primären Phloems bei *Cucurbita ficifolia*. Planta 59:243-61.
1963b. Der Phloemsaft von *Cucurbita ficifolia*. Planta 60:216-24.
- EVERT, R. F.
1963. The cambium and seasonal development of the phloem in *Pyrus malus*. Amer. Jour. Bot. 50:149-59.
- EVERT, R. F., and W. F. DERR.
1964a. Callose substance in sieve elements. Amer. Jour. Bot. 51:552-59.
1964b. Slime substance and strands in sieve elements. Amer. Jour. Bot. 51:875-80.
- EVERT, R. F., and L. MURMANIS
1965. Ultrastructure of the secondary phloem of *Tilia americana*. Amer. Jour. Bot. 52:95-106.
- FRIEDMAN, I., and E. S. BIRD
1960. Pleomorphic cytoplasmic inclusion bodies in tissue cultures of the otocyst exposed to dimycin. In: Fourth International Conference on Electron Microscopy, 1958, pp. 266-70. Springer-Verlag, Berlin.
- GENÈVES, L.
1964. Origine des espaces intercellulaires dans les méristèmes radiculaires de *Raphanus sativus* L. (Crucifères). Acad. des Sci. Compt. Rend. 258:302-04.
- HEPTON, C. E. L., and R. D. PRESTON
1960. Electron microscopic observations of the structure of sieve-connexions in the phloem of angiosperms and gymnosperms. Jour. Exptl. Bot. 11:381-93.
- HILL, A. W.
1908. The histology of the sieve-tubes of angiosperms. Ann. Bot. 22:245-90.
- ITO, S.
1962. Post-mortem changes of the plasma membrane. Fifth International Congress for Electron Microscopy. Vol. 2, p. L-5. New York: Academic Press.

- KESSLER, G.
1958. Zur Charakterisierung der Siebröhrenkallose. Schweiz. Bot. Gesell. Ber. 68:5-43.
- KOLLMANN, R.
1960a. Untersuchungen über das Protoplasma der Siebröhren von *Passiflora coerulea*. I. Mitteilung. Lichtoptische Untersuchungen. Planta 54:611-40.
1960b. Untersuchungen über das Protoplasma der Siebröhren von *Passiflora coerulea*. II. Mitteilung. Elektronenmikroskopische Untersuchungen. Planta 55:67-107.
- KOLLMANN, R., and W. SCHUMACHER
1964. Über die Feinstruktur des Phloems von *Metasequoia glyptostroboides* und seine jahreszeitlichen Veränderungen. V. Mitteilung. Die Differenzierung der Siebzellen im Verlaufe einer Vegetationsperiode. Planta 63:155-90.
- KÜSTER, E.
1951. Über die Siebplatten und Siebfelder der *Cucurbita*-Siebröhren und ihren Kallus. Ztschr. f. Mikros. 60:157-62.
- MANTON, I.
1961. Some problems of mitochondrial growth. Jour. Exptl. Bot. 12:421-29.
1962. Observations on stellate vacuoles in the meristem of *Anthoceros*. Jour. Exptl. Bot. 13:161-67.
- MEHTA, A. S., and D. C. SPANNER
1962. The fine structure of the sieve tubes of the petiole of *Nymphoides peltatum* (Gmel.) O. Kunze. Ann. Bot. 26:291-99.
- MENKE, W.
1959. Myelinfiguren im Cytoplasma meristematischer Zellen. Protoplasma 51:127-30.
- METCALFE, C. R., and L. CHALK
1950. Anatomy of the dicotyledons. Vol. 1. Oxford: Clarendon Press.
- PAOLILLO, D. J., JR.
1963. The developmental anatomy of *Isoetes*. Illinois Biological Monographs 31. Urbana: The University of Illinois Press.
- ROBERTSON, J. D.
1960. A molecular theory of cell membrane structure. In: Fourth International Conference on Electron Microscopy, 1958, pp. 159-71. Springer-Verlag, Berlin.
- SCHANDERL, H.
1957. Über das Vorkommen und die Bildungsweise der Gerbstoffe und der Anthozyane in den Rebengewächsen. Rebe und Wein, Mitt. Ser. A. 7:229-45.
- SINNOTT, E. W., and R. BLOCH
1941. Division in vacuolate plant cells. Amer. Jour. Bot. 28:225-32.
- THAINE, R.
1964. Protoplast structure in sieve tube elements. New Phytol. 63:236-43.
- WARDROP, A. B., and J. CRONSHAW
1962. Formation of phenolic substances in the ray parenchyma of angiosperms. Nature 193:90-92.
- WHALEY, W. G., H. H. MOLLENHAUER, and J. H. LEECH
1960. The ultrastructure of the meristematic cell. Amer. Jour. Bot. 47:401-49.
- WILHELM, K.
1880. Beiträge zur Kenntnis des Siebröhrenapparates dicotyler Pflanzen. Wilhelm Engelmann, Leipzig.
- WOODING, F. B. P., and D. H. NORTHCOTE
1965. The fine structure and development of the companion cell of the phloem of *Acer pseudoplatanus*. Jour. Cell Biol. 24:117-28.
- ZIRKLE, C.
1932. Vacuoles in primary meristems. Ztschr. f. Zellforsch. u. Mikros. Anat. 16:26-47.

To simplify the information, it is sometimes necessary to use trade names of products or equipment. No endorsement of named products is intended nor is criticism implied of similar products not mentioned.

The journal HILGARDIA is published at irregular intervals, in volumes of about 650 to 700 pages. The number of issues per volume varies.

Single copies of any issue may be obtained free, as long as the supply lasts; please request by volume and issue number from:

**Agricultural Publications
University Hall
University of California
Berkeley, California 94720**

The limit to nonresidents of California is 10 separate titles. The limit to California residents is 20 separate titles.

The journal will be sent regularly to libraries, schools, or institutions in one of the following ways:

- 1. In exchange for similar published material on research.**
- 2. As a gift to qualified repository libraries only.**
- 3. On a subscription basis—\$7.50 a year paid in advance. All subscriptions will be started with the first number issued during a calendar year. Subscribers starting during any given year will be sent back numbers to the first of that year and will be billed for the ensuing year the following January. Make checks or money orders payable to The Regents of The University of California; send payment with order to Agricultural Publications at above address.**