HILGARDIA

A JOURNAL OF AGRICULTURAL SCIENCE PUBLISHED BY
THE CALIFORNIA AGRICULTURAL EXPERIMENT STATION

Volume 37, Number 4 • December, 1965

Translocation Relationships in and between Mistletoes and Their Hosts

O. A. Leonard and R. J. Hull

UNIVERSITY OF CALIFORNIA DIVISION OF AGRICULTURAL SCIENCES

Studies of translocation between mistletoes and their host trees were conducted under natural conditions in various California locations. Both dwarf (*Arceuthobium* spp.) and green (*Phoradendron* spp.) mistletoes were studied. Various labeled materials were used, including $C^{14}O_2$, several C^{14} -labeled herbicides, and P^{32} and S^{35} (as phosphate and sulfate, respectively).

The C^{14} -labeled assimilates and the P^{32} translocated from host foliage into dwarf mistletoes growing on the host branches, but no such movement occurred from host branches to the mistletoes. When $P^{32}O_4$ was applied to the wood of the host branches, beneath the infections, P^{32} appeared in both dwarf and green mistletoes. This type of application apparently was a highly effective means of introducing the labeled phosphate.

Labeled substances applied to the shoots of dwarf mistletoes normally moved apically. In a few instances, P^{32} was found to move basally into the endophytic system and host branch. This movement was not through living tissues, however, because it was not blocked by killing the basal parts of the mistletoe shoots with steam.

Translocation in green mistletoes differed from that in dwarf mistletoes. In the green species, labeled assimilates and P^{32} were transported basipetally into the endophytic systems. However, label did not move from the green mistletoes into the host branches.

THE AUTHORS:

O. A. Leonard is Lecturer in Botany and Botanist in the Experiment Station, Davis. R. J. Hull was formerly Research Assistant in the Department of Botany, Davis; now with the Department of Botany and Plant Pathology, Purdue University, Lafayette, Indiana.

Translocation Relationships in and between Mistletoes and Their Hosts¹

INTRODUCTION

MISTLETOES in the United States are represented by two genera: the light-green or dwarf mistletoes (*Arceuthobium* spp.) that grow on some coniferous species; and the green mistletoes (*Phoradendron* spp.) that grow on both coniferous and dicotyledonous woody plants. Both are destructive, parasitic flowering plants. That they are of serious economic importance was shown in a review by Gill and Hawksworth (1961). Offord (1960) had previously noted that dwarf mistletoes are among the most important forest parasites of western North America.

To date, progress on chemical control of mistletoes is limited, as are studies on translocation of substances between them and their hosts. In the work reported here, translocation patterns between mistletoes and their hosts were studied by means of several labeled materials, including $C^{14}O_2$ (for producing labeled assimilates), several C^{14} -labeled herbicides, and P^{32} and S^{35} (as phosphate and sulfate, respectively). We hoped to learn whether characteristic patterns of transport existed between dwarf and green mistletoes and

their respective hosts, and whether such patterns might differ from the normal. Transport patterns in noninfected host plants were also studied to determine whether mistletoes influenced normal translocation. Surveys were made of the effect of method and point of application of labeled material on transport. Certain aspects of these studies have been presented (Hull and Leonard, 1964a,b), including qualitative and quantitative data on the carbohydrates translocated from labeled assimilates. The present paper supplements those data, reports results obtained from all the autoradiographic studies (about 300), and indicates aspects of translocation not evident in quantitative analyses of whole plant parts. It is hoped that translocation data will prove helpful in developing chemical controls for dwarf and green mistletoes. Such data should also be useful in other areas of research, including the use of systemic plant protectants, nutrients, or herbicides.

Except for a few greenhouse tests, all studies were made in the field, under natural conditions, and results should be applicable under those conditions.

REVIEW OF LITERATURE

Only work relating to translocation and control will be discussed. The review by Gill and Hawksworth in 1961 contained what was known to that

time on the translocation relationship between mistletoes and hosts, and will not be repeated here except where it is considered relevant.

¹ Submitted for publication November 13, 1964.

Translocation

Hull and Leonard (1964a,b) observed that host foliage supplied carbohydrates to the dwarf but not to the green mistletoes. There was no indication that mistletoe infections blocked the flow of assimilates transported by phloem; however, the dwarf mistletoe did act as a physiological block at times by diverting much of the available food from the host to the parasite. Hull and Leonard further observed that carbon fixed by the dwarf and green mistletoes did not enter the hosts. This result contrasts with the finding of Rediske and Shea (1961) that assimilates from *Arceuthobium americanum* on lodgepole pine (*Pinus murrayana*) did actually enter the host from the parasite. It is suggested, however, that their finding resulted from leakage of a small amount of labeled CO₂ which was transformed into assimilates by the pine leaves.

Wagener (1925) observed that a green mistletoe (*Phoradendron juniperinum* var. *libocedri*) on incense cedar (*Libocedrus decurrens*) had remained alive for more than 100 years, even though it was visibly without shoots. This observation suggests that small quantities of food materials might be obtainable from the host. Further studies are necessary to determine whether this in fact occurs in sufficient quantities to sustain life.

On the basis of field observations, Weir (1916) suggested that dwarf mistletoes can supply assimilates to defoliated hosts, but the findings of Hull and Leonard (1964b) indicate that this is unlikely. Several authors have expressed similar ideas with respect to the green mistletoes supplying food to defoliated host branches (Gill and Hawksworth, 1961). Definite data on this question are supplied in the present paper.

Greenham and Brown (1957), working in Australia, applied Fe⁵⁹Cl₃ to seven holes in a 7.2-inch *Eucalyptus dives* trunk infected in two areas by *Amyema pendula*. The eucalyptus leaves became variably labeled within a few days, since

certain branches received much more label than did others. During this period, the mistletoe remained free of the label. After one year, the host leaves became more uniformly labeled, and both mistletoes had about 50 per cent as much label as did the host foliage. Thus, considerable redistribution of the label occurred with time.

Greenham and Leonard (1964) observed considerable similarity between the amino acids in mistletoes and those in the hosts. However, certain free amino acids present in the mistletoes were not present in the host tissues. This information suggested that amino acids might pass from hosts into mistletoes but that the reverse was not indicated. Some possibility of control was suggested by applying analogs of the amino acids not common to both host and parasite. Possibilities of this kind are suggested by the remarkable specificity of forms of some mistletoes for their hosts. For example, *Arceuthobium campylopodum* on *Abies concolor* (white fir) does not attack *A. magnifica* (red fir), and the same species of dwarf mistletoe on red fir does not attack white fir (Parmer and Scharpf, 1963).

Chemical Control

Hartigan (1949) was able to effect partial control of *Amyema* on eucalyptus by spraying the mistletoes with 2,4-dichlorophenoxyacetic acid (2,4-D). Unfortunately, the sprays frequently resulted in an unacceptable amount of damage to the eucalyptus, which is evergreen, and 2,4-D has therefore never been used on a large scale. Graser (1954) made similar observations on *Phoradendron* control on walnut (*Juglans*), using 2,4-D amine. Control of the majority of infections by means of dormant applications was possible, without injury to the walnut, when the host was free of leaves. The method is tedious because the 2,4-D must be applied only to the mistletoes. Spraying has not resulted in control of dwarf mistletoes except when the infections themselves were

TABLE 1
MISTLETOES AND HOST TREES STUDIED IN CALIFORNIA,
TEST SITES, AND ELEVATION

Mistletoe	Host trees	Test site	Elevation
			meters
<i>Arceuthobium</i> (dwarf):			
<i>americanum</i>	lodgepole pine (<i>Pinus murrayana</i>)	Hermit Valley	2,150
<i>campylopodum</i>	Jeffrey pine (<i>P. jeffreyi</i>)	Cow Creek	1,890
	ponderosa pine (<i>P. ponderosa</i>)	Strawberry	1,675
	sugar pine (<i>P. lambertiana</i>)	Strawberry	1,770
	single leaf pinyon (<i>P. monophylla</i>)	Walker	1,650
	digger pine (<i>P. sabiniana</i>)	Knoxville	400
		Mount Diablo	760
	white fir (<i>Abies concolor</i>)	Strawberry	1,770
	red fir (<i>A. magnifica</i>)	Pinecrest	2,225
<i>Phoradendron</i> (green):			
<i>bolleanum</i> var. <i>densum</i>	Macnab cypress (<i>Cupressus macnabiana</i>)	Knoxville	760
<i>bolleanum</i> var. <i>pauciflorum</i>	white fir (<i>Abies concolor</i>)	Strawberry	1,700
<i>flavescens</i> var. <i>macrophyllum</i>	California black walnut (<i>Juglans hindsii</i>)	Davis	16
<i>flavescens</i> var. <i>villosum</i>	blue oak (<i>Quercus douglasii</i>)	Knoxville	450
	California white oak (<i>Q. lobata</i>)		
	interior liveoak (<i>Q. wislizenii</i>)	Hopland	760
	California black oak (<i>Q. kelloggii</i>)	Hopland	760
<i>juniperinum</i> var. <i>libocedri</i>	incense cedar (<i>Libocedrus decurrens</i>)	Strawberry	1,700
<i>juniperinum</i> var. <i>ligatum</i>	western juniper (<i>Juniperus occidentalis</i>)	Eagle Meadow	2,200

treated (Quick, 1964). The most promising herbicide appears to be the isooctyl ester of 4(2,4,5-trichlorophenoxy) butyric acid (2,4,5-TB), but this will be applicable only for very high-value trees because complete coverage of each infection is required—a time-consuming and difficult operation, especially on large trees.

Trunk injection with 2,4-D and 4(2-methyl-4-chlorophenoxy) butyric acid (MCPB) is being used in Australia to control mistletoe on eucalyptus infected with *Amyema pendula* (Greenham and Brown, 1957; Brown, 1959; Greenham, 1964). This is the most widely used chemical method of mistletoe control at present, and even its use is limited because of the labor involved in making the applications. Kadambi (1954) found

that copper sulfate injected into hosts controlled *Loranthus pulverulentus*.

Apparently the selective control of mistletoes is not insurmountable provided economical and sufficiently selective materials can be found. This is suggested by the work of Moss (1960) in treating white pine blister rust (*Cronartium ribicola*) on western white pine (*Pinus monticola*) with cycloheximide (Acti-dione). The antibiotic was applied in fuel oil to the trunks of the trees, from which it was gradually absorbed and translocated to the cankers. Aircraft spraying of another antibiotic (phytoactin) is being tested in large-scale field operations (Viche, Moss, and Hartmann, 1962; Anonymous, 1962). Perhaps someday we shall see similar use of sprays for controlling mistletoes.

MATERIALS AND METHODS

Nearly all of the studies were conducted on 15- to 50-year-old trees growing naturally in forests of the Sierra Nevada and Coast Range. A list of the hosts and mistletoes studied and the location and elevation of the test sites are given in table 1. The greatest empha-

sis in the present study was on the dwarf mistletoes (*Arceuthobium* spp.), but sufficient work was done on the green mistletoes (*Phoradendron* spp.) to establish the translocation patterns within those plants and between them and their hosts.

Fig. 1. Apparatus used to expose terminal foliage of mistletoe-infected branches to $C^{14}O_2$: cg, carbon dioxide generator; cp, circulatory pump; dm, dwarf mistletoe shoots; pb, polyethylene bag.

In one study, the movement of labeled assimilates in and between both mistletoes and their hosts was determined. The labeled assimilates were produced in sunlight from $C^{14}O_2$ ($50 \mu\text{c}$ per treatment) following procedures described by Hull and Leonard (1964*a,b*) and illustrated in figures 1 and 2. The polyethylene bags containing the $C^{14}O_2$ were kept on the branches about four hours. Heat damage was avoided by applying treatment in the forenoon and by partial shading of the branches when considered necessary.

Other studies were conducted with P^{32} and S^{35} (as the sodium or ammo-

nium salts of phosphate and sulfate) adjusted to a pH of 6 to 7. About $100 \mu\text{c}$ of radioactivity were applied, per treatment, either to host foliage or wood, or to mistletoe shoots. In some cases the mistletoe shoots were pricked with a pin to facilitate uptake of the materials. In some tests the mistletoe shoots were steamed at the base in order to study the effect of killing that portion of the shoot on the translocation of label through the steamed portion. Death of the steamed portion was determined by means of a resistance bridge described by DePlater and Greenham (1959). The treated areas were protected from rain

Fig. 2. Apparatus used to expose individual shoots of dwarf mistletoe to $C^{14}O_2$ without detaching them from their host plant.

by polyethylene bags which were open on the undersides. Additional aspects of the various treatments are described under specific experiments.

The following compounds were applied in 50 per cent ethanol containing 0.1 per cent Tween 20 (polyoxyethylene sorbitan monolaurate): C^{14} -labeled urea (specific activity (s.a.) 6.2 mc/mmole); 2,4-D (carboxyl labeled with s.a. of 1.24 mc/mmole); 2,4,5-T (s.a. 1.0 mc/mmole); butoxy ethanol ester of 2,4,5-T (s.a. 0.94 mc/mmole); 2-chloro-4-ethylamino-6-isopropylamino-s-triazine (atrazine; s.a. 1.4 mc/mmole, randomly ring-labeled); 3-amino-1,2,4-

triazole-5- C^{14} (amitrole; s.a. 0.94 mc/mmole); paraquat (1,1'-dimethyl-4,4'-dipyridylum cation; s. a. 1.44 mc/mmole); and maleic hydrazide (MH; s.a. 1.2 mc/mmole). One μ c of activity, in 10 to 30 μ l, was applied either to the host foliage, bark, or wood, or to the mistletoe shoots.

Treated branches were usually removed from the trees two weeks after treatment. (Shorter time intervals proved inadequate in several tests; part of these data is shown in table 5.) Stem sections of host were cut either with a razor blade or a band-saw to a thickness of 1 to 2 mm. The distance between sec-

tions from which cuts were made was noted on the mounting paper. Portions of the branch and the infecting mistletoe were mounted on 10 × 12-inch sheets of paper by means of a casein-base glue. Mounts were pressed between wire screens and dried at 45° to 50° C. Dried mounts were rehumidified and pressed in a mechanical press so that the surface to be exposed to the film was as smooth as possible. In a few cases the mounts were freeze-dried. The autoradiographic procedure was similar to that of Yamaguchi and Crafts (1958), except that a method of preventing pseudoautoradiographs was used. These are produced on X-ray film by natural chemicals in the plant materials, not by radiation. We found that they could be prevented (except for some faint images at times—for example, fig. 3) by certain film materials. Saran Wrap (Dow Chemical Company) was chosen because it eliminated most pseudoautoradiographs while transmitting over 50 per cent of the beta radiation. Autoradiographs were made by placing Saran Wrap-

covered mounts on Kodak Royal Blue Medical X-ray film for two weeks.

The distribution of radioactivity in the autoradiographs was estimated by comparing the densities with a series of standards of known activity (counts per minute per mg), corrected for self-absorption. The series and standards (cpm/mg) used were: 0 (no definite activity); 1 (0–0.5); 2 (0.6–2); 3 (3–9); 4 (9–32); 5 (33–128); 6 (129–712); 7 (more than 712). Estimates of radioactivity for all host stem cross-sections are for the inner portions of the bark and adjacent cambial area. Variations in radioactivity are listed for the sections chosen—for example, a given cross-section may be free of activity on one side (0) and very radioactive on the other (7), and thus be listed at 0–7. Different mistletoe shoots varied in activity, and such variation is noted.

A Beckman L/B infrared gas analyzer was used to study respiration and photosynthesis of *Phoradendron* by the same procedure as that of Hull and Leonard (1964b) with *Arceuthobium*.

RESULTS

DWARF MISTLETOES

Basipetal Translocation of C¹⁴ in Conifer Branches

White fir (*Abies concolor*), red fir (*A. magnifica*), digger pine (*Pinus sabiniana*), sugar pine (*P. lambertiana*), Jeffrey pine (*P. jeffreyi*), ponderosa pine (*P. ponderosa*), lodgepole pine (*P. murrayana*), and pinyon pine (*P. monophylla*) were treated during the growing season, and all except lodgepole, pinyon, and sugar pine were also treated while dormant (tables 2, 3, and 4; figs. 4, 5, 6, and 7). Considerable variation appeared in the labeling of the stem cross-sections of the conifers and the endophytic systems and aerial shoots of the dwarf mistletoes. Labeling was uniform when no untreated side branches were present between the

treated area and the tissues sampled. The effectiveness of labeling became decidedly nonuniform at points where the treated branch joined large untreated branches. Some lateral mixing of the translocation streams did occur since all portions of the endophytic systems, aerial dwarf mistletoe shoots, and stem cross-sections usually had at least a trace of label; however, in some instances certain parts of the cambial area of the stem cross-sections appeared to be completely free of label.

Although assimilates were translocated out of the treated area during the two-week treatment period at all seasons of the year, large quantities of labeled carbon always remained within the treated area. Translocation was most

Fig. 3. Effectiveness of Saran Wrap in eliminating pseudoautoradiographs of digger pine and Macnab cypress. A. mounted branch; B. pseudoautoradiograph produced by exposing X-ray film to mounted branch for two weeks; C. pseudoautoradiograph produced by exposing X-ray film for two weeks to mounted branch covered with Saran Wrap.

rapid before the shoots began to elongate in the spring (table 5, for February and June). Translocation became slow in white fir when the shoots were rapidly elongating in July (quite variable) and August (tables 2 and 5). (See Hull and Leonard, 1964a, for carbohydrate studies.) During that period, the autoradiographs showed intense activity in the new terminal growth of the shoots. In one instance, no labelling appeared in the endophytic system and aerial shoots of the dwarf mistletoes even though the label had traversed that area to the base of the stem (table 2, July, 1961, in the early active-growth period). In two instances, however, the assimilates had reached and were accumulated by the dwarf mistletoes without moving beyond to the base of the stems in white fir (table 2, July, 1961 and 1962). Simi-

lar results were observed in sugar, Jeffrey, ponderosa, and pinyon pine, and red fir, when the shoots were elongating (table 3), and in digger pine after shoot growth had ceased (table 3, August, 1961). Downward translocation in white fir still appeared to be very slow in October after all shoot growth had ended (fig. 7 and table 5).

While downward translocation generally appeared to be similar in infected and noninfected stems (fig. 7 and table 5), indications were that the infections did at times interfere with transport (fig. 5). For example, photosynthate never failed to reach the base of treated, noninfected stems within two weeks, but such failure did occur in two instances out of 150, with infected white fir stems (table 2, July, 1961 and 1962). Additional instances of assimilates not mi-

grating beyond the infected areas in several conifers (table 3) strongly suggest that the infections blocked translocation, probably by using all of the food not used by the shoot itself during the summer. However, large infections on digger pine (fig. 4E), ponderosa pine, Jeffrey pine, and red fir (fig. 4C) failed to block the transport of assimilates (table 3). Thus dwarf mistletoe infections apparently do not completely block translocation except occasionally when food requirements of both the hosts and the dwarf mistletoes are high, and coincide.

The endophytic system was a strong sink for assimilates coming from the coniferous hosts at all seasons of the year, while the aerial shoots were strong sinks during the summer period (tables 2 and 3), and somewhat less so at other times of the year. However, the latter always accumulated appreciable assimilates from the host.

An experiment was initiated in 1960 on white fir to determine whether dwarf mistletoe shoots and/or exposure of the infections to light was necessary for the accumulation of assimilates by the endophytic systems. Dwarf mistletoe shoots were removed from some branches; other infected branches were left with shoots intact. Half the branches from which shoots were removed were covered with aluminum foil at the site of the infection, to keep out light. Infections with shoots intact were also covered with the foil. The host foliage was treated with $C^{14}O_2$ at various periods following preparation, and the accumulation of assimilates in the host and mistletoe was noted in the autoradiographs (table 4 and fig. 6). Removing the shoots appeared to increase the accumulation of label in the endophytic system, but covering the infections with aluminum foil had little effect. Some of the covered branches selected for treatment in July, 1960, died in eight to 12 months; however, considerable mortality is common with white fir branches infected with dwarf mistletoe.

Wood in the treated shoots of conifers frequently contained appreciable label in the xylem rays and resin ducts. Storage of assimilates in the rays was apparently greatest before growth started in spring and during the summer, and least during the winter. Large quantities of label were present in the resin ducts of digger, lodgepole, and pinyon pine just before, during, and just after the period of rapid shoot growth.

Effect of Host Defoliation on Translocation of C^{14} in Noninfected and Infected Branches

Data in table 6 and figures 8 and 9A show the influence of defoliation on translocation in healthy and infected white fir lateral branches. Movement into the defoliated, infected branches was extensive at all times of the year. Movement into aerial shoots was greatest during the growing season. Translocation into uninfected, defoliated branches was intense only during the growing season.

Intense translocation into nondefoliated, infected lateral branches did occur in a few instances. Assimilates moved into infected white fir shoots in the early active period of shoot growth (table 6 and fig. 6C) and into a similar lodgepole pine shoot at the same stage of growth. Limited upward movement (for 2 cm) in an infected white fir branch occurred in November, 1960 (fig. 8C).

Defoliation of noninfected branches during the growing season induced import of large quantities of assimilates, while defoliation at other seasons of the year was far less effective in bringing this about (table 6). Covering the branches with aluminum foil was less effective than defoliation in inducing branches to import assimilates (fig. 9B).

No instance was observed in which noninfected, nondefoliated, normal branches systemically imported label from treated branches; however, limited movement into the basal 1 to 6 cm was

TABLE 2

SELECTED SAMPLES SHOWING DISTRIBUTION OF C¹⁴-LABELED ASSIMILATES
IN WHITE FIR BRANCHES INFECTED WITH DWARF MISTLETOE
(Radioactivity determined two weeks after labeling of terminal foliage)

Growth stage of host, and date of labeling	Intensity of autoradiographs*			
	White fir stem		Mistletoe	
	Area above infection	Base of stem	Endophytic system	Aerial shoots
Dormant:				
1/29/63.....	0-7	0-4	...	...
1/29/63.....	0-7	1-5	2-7	2-3
2/5/61.....	1-7	2-7	1-7	0-1
2/7/61.....	7	1-7	7	2
2/7/63.....	1-7	1-7	...	...
2/14/63.....	7	1-5	2-7	2-4
2/15/63.....	2-7	2-7	...	...
5/4/63.....	7	2-7	3-7	2-5
5/30/61.....	7	7	...	...
5/30/61.....	7	2-7	2-7	1-2
5/30/62.....	7	5-7	...	...
6/4/63.....	7	2-6	...	...
6/4/63.....	7	1-7	2-7	1-2
6/4/63.....	2-7	2-5	1-7	2-5
6/5/64.....	7	0-6	2-7	4-6
6/13/62†.....	7	2-6	2-7	3-5
6/30/63†.....	7	1-7	2-7	1-3
6/30/62†.....	7	3-7	...	...
Early-active:				
7/3/61.....	7	0-2	0	0
7/3/61.....	7	3	7	7
7/5/64.....	7	0-5	0-5	2-5
7/5/64.....	7	1	0-4	2-4
7/18/62.....	7	1-3	...	...
7/18/62.....	7	0	1-7	6-7
7/25/61.....	7	0	0-7	1-7
7/25/61.....	7	0-5	...	...
Active and postactive:				
8/7/61.....	1-7	0-5	1-7	1-6
8/16/60.....	7	1	...	...
8/16/60.....	7	1-5	7	7
8/31/62.....	7	2-7	7	7
9/3/62.....	7	0-6	7	6
9/3/62.....	7	0-6	...	...
9/12/63.....	4-7	0-5	...	...
Dormant:				
10/2/62.....	7	2-5	7	1-5
10/2/62.....	6	2-5	...	...
11/1/61.....	6	0-5	6	3-5
11/20/60.....	2-7	0-2	7	1

* Radioactivity ratings are on a scale from 0 to 7, on the basis of counts per minute per mg. For a fuller explanation see text, p. 118.

† Dormant to bud-swell.

observed in 12 branches. This movement mainly occurred either shortly before or during start of growth in spring.

There was evidence of xylem transport from treated to untreated branches of white fir just before bud-break in the spring (fig. 7B, C). Leaves of untreated, infected and noninfected branches were lightly labeled; no label accumulated in the cambial area of such shoots, but did accumulate in the treated branches.

Effect of Girdling on Downward Translocation in Infected Branches

Initially it was thought that dwarf mistletoe infections might produce effects similar to those caused by certain types of tree girdling. To investigate

TABLE 3

SELECTED SAMPLES SHOWING DISTRIBUTION OF C¹⁴-LABELED ASSIMILATES IN BRANCHES OF VARIOUS CONIFERS INFECTED WITH DWARF MISTLETOE (Radioactivity determined two weeks after labeling of terminal foliage)

Host and stage of growth	Date of labeling	Intensity of autoradiographs*			
		Host stem		Mistletoe	
		Area above infection	Base of stem	Endophytic system	Aerial shoots
Digger pine:					
Dormant.....	3/23/61	7	5-7	7	4-5
	11/1/61	6	0-4	6	3-5
Active.....	5/11/62 large infection	7	7	7	7
	7/10/62	7	7	7	7
Post-active.....	8/22/61	7	0-4	7	1-7
	8/29/61	7	0	7	4-5
Lodgepole pine:					
Active.....	7/18/62	7	1-5	7	7
	8/10/62	7	1-7	1-7	2-7
Sugar pine:					
Active.....	7/25/61	7	0	1-7	1-7
Ponderosa pine:					
Dormant†.....	6/5/64	7	0-6	7	6-7
	6/5/64 large infection	7	0-4	5	7
Active.....	8/22/61	7	0	1-7	1-7
Jeffrey pine:					
Dormant†.....	6/5/64	7	0-6	7	5-7
	6/5/64	5	0-4	2-6	2-5
Active.....	8/8/61 large infection	5	0	0-7	3-7
Pinyon pine:					
Active.....	8/10/61	7	0-4	1-7	3-4
	8/10/61	7	0	1-7	2-6
Red fir:					
Dormant.....	6/5/64	7	0-7	2-7	2-6
	6/5/64 large infection	5	2-5	1-5	1-3
Active.....	7/25/61	7	0	2-7	2-7

* Radioactivity ratings are on a scale from 0 to 7, on the basis of counts per minute per mg. For a fuller explanation, see text, p. 118.

† Dormant to bud-swell.

this possibility, we studied the effect of several types of girdles on translocation (table 7). Three out of five white fir branches with simple (complete) girdles did not translocate assimilates to the girdles. In one instance there was no basal translocation out of the treated area, but there was strong movement into the new growth at the tips (July, 1963, when the shoots were rapidly growing). Translocation to the girdle occurred when the girdle had been made by killing the bark with fire. Transport in a detached branch, which represents a type of girdling, was similar to that obtained by a simple girdle (July,

1960). Double-spiral and loop girdles (partial girdles) had a continuous strip of bark connecting the undisturbed bark above and below the portion of the stem from which the bark was removed. The assimilates were able to move laterally in and through the double spiral to the bark below. In one instance, movement did not proceed farther than 8 cm below the spiral, while in another, movement did proceed to the base of the stem. With the loop girdle, the strip of continuous bark was cut so that materials moving in the bark had to move upward for 6 cm before moving downward again. Assimilates were able to move

Fig. 5. Autoradiographs showing no translocation beyond infected areas of conifer branches infected with *Arceuthobium campylopodum*. Terminal foliage (above arrows) was exposed to $C^{14}O_2$, and collections were made two weeks later. The X marks location of lowest stem cross-sections. In all cases, mistletoe infection was actively growing and producing staminate or pistillate flowers and, in most cases, fruit. Species treated as follows: A. sugar pine, July 25, 1961, during active growth; B. ponderosa pine, August 22, 1961, just following active growth; C. red fir, July 25, 1961, during active growth; D. digger pine, August 29, 1961, after active growth had stopped; E. Jeffrey pine, August 8, 1961, during active growth; F. white fir, July 18, 1962, during active growth.

through the loop and accumulate in the infected area below.

Assimilates were transported to a simple girdle in lodgepole pine in high concentration, and were detectable in the pith to 10 cm beyond the girdle (August, 1962). Translocation of assimilates in digger pine was interrupted by the simple girdles, but label was present in the resin canals for 15 cm below the girdles. Assimilates were able to move through a double spiral of bark and into the dwarf mistletoe, where they accumulated.

The downward movement of $P^{32}O_4$ in digger pine was similar to that with assimilates. However, $P^{32}O_4$ was not so

confined by a girdle as were assimilates, and was frequently present for several inches beneath the girdles. In one instance the $P^{32}O_4$ evidently leaked into the xylem, for the label was transported down and then up a lateral branch, through a girdle.

Translocation of C^{14} Following Treatment of Mistletoe Infections

Dwarf mistletoe infections (host tissues plus the associated mistletoe endophytic system and aerial shoots) on white fir branches were treated with $C^{14}O_2$. That portion of the host stem exposed to the $C^{14}O_2$ was free of leaves.

Fig. 6. Mounted plants (above) and autoradiographs (below) of *Arceuthobium campylopodium*-infected white fir branches, terminal foliage of which (above arrow) was exposed to $C^{14}O_2$. Collections were made two weeks later. Dwarf mistletoe shoots were removed seven months or longer before treatments. Note large accumulation of label in infected area (es). A. treatment February 5, 1961. Infected area was covered with aluminum foil. B. treatment May 30, 1961. Infected area was covered with aluminum foil. C. treatment July 3, 1961. Note accumulation of label by young fir shoot in infected area (es).

The treatments were made at five periods of the year, and included all seasons.

Collections were made two weeks after the $C^{14}O_2$ was applied. About 60 times as much label was present in the region of infection (endophytic system plus associated host tissues) from host foliage treatments as from treatment of the entire dwarf mistletoe (plus associated host branch tissues). Slightly more label appeared in the infected branches than in similarly treated, non-infected branches; this may have been due to the occurrence of more breaks in the host bark in the infected areas. Label was present for about 10 cm both above and below the portion of the branches (infected and noninfected)

exposed to $C^{14}O_2$ and most of the label was present in the outermost part of the bark. This finding contrasted with the high concentration of label in the inner part of the bark of the host and the mistletoe endophytic system when the host foliage was treated with $C^{14}O_2$. The mistletoe shoots exposed to $C^{14}O_2$ contained about 35 per cent as much label (in two weeks) as did those that imported label coming from the host foliage.

Translocation of C^{14} and Labeled Phosphate Following Treatment of Mistletoe Shoots

In order to eliminate uptake of label by the host bark as a factor influencing the results, only the mistletoe shoots

Fig. 7. Autoradiographs of *Arcuthobium campylopodum*-infected white fir branches, terminal foliage of which (above arrow) was exposed to $C^{14}O_2$ for four hours. Tests were conducted at different periods of the year to determine seasonal differences in rates of translocation. Infected area (endophytic system) is at (E). Dates and time after treatment: A. January 29, 1963, 28 hours; B. June 3, 1963, 10 hours; C. June 3, 1963, 47 hours; D. August 7, 1961, 34 hours; E. August 7, 1961, 57 hours; F. October 2, 1961, 52 hours.

were treated with $C^{14}O_2$ (fig. 2). Only that portion of the shoot actually exposed to $C^{14}O_2$ contained any label (table 8 and fig. 10). No basal movement of label occurred in the mistletoe shoots in any test. (Many tests were conducted on white fir, digger pine, and lodgepole pine.) The label was concentrated in the outermost part of the mistletoe stems, at the bases of the internodes, in the male and female flowers, and in the fruit. Mistletoe shoots that received label from the host foliage were labeled in about the same manner as described above. The inner portions of the shoots were more strongly labeled, however, and that portion containing the xylem (fig. 18B) was nearly free of label.

Dwarf mistletoe shoots treated with $P^{32}O_4$ usually did not show any label moving into the endophytic system or into the host tissues; under some conditions, however, puncturing the surfaces in several places with a pin, before treatment, did result in movement of a considerable amount of label into the endophytic system and into the host (table 8 and fig. 11). Natural breaks in the tissues would also result in some label moving from the aerial shoots into the endophytic system and host tissues; this occurred in four of the trials shown in table 8. Puncturing the shoots in six to eight places resulted in movement of $P^{32}O_4$ into the endophytic system during the late fall and winter period, but was ineffective in late spring.

TABLE 5
TRANSLOCATION RATES OF C¹⁴-LABELED ASSIMILATES, AT DIFFERENT SEASONS OF THE YEAR, IN NORMAL BRANCHES OF WHITE FIR AND IN BRANCHES INFECTED WITH DWARF MISTLETOE

Growth stage of host	Month treated	Translocation* of C ¹⁴ in:	
		Infected branches	Noninfected branches
		cm/hr.	cm/hr.
Dormant.....	January	2.1,† 2.1†	2.7†
Dormant.....	February	2.4,† 2.1†	2.7†
Dormant.....	June	2.6, 3.0, 6.0, 2.0†	5.0, 2.5
Active.....	July	0.2, 1.8, 1.8, 0.0	2.0, 1.0
Active.....	August	0.5, 0.3, 0.0, 1.0, 0.5	0.7, 0.2, 0.3
Dormant.....	October	0.2, 0.0, 0.2, 0.2, 0.3	0.0, 0.0, 0.3, 1.0

* Translocation rates were determined by noting the branch cross-sections that were radioactive (as shown by autoradiographs). Collections were made from seven hours to 14 days following labeling of terminal foliage.
† Translocation rates were in excess of those indicated, since C¹⁴ was already at the base of the branches when collections were made.

In three out of six trials in which the bases of the mistletoe shoots were killed with steam before application of P³²O₄, appreciable label appeared in the endophytic system. Since the bases were killed before application of the label, the movement had to occur from the shoots to the endophytic system through dead tissues. (The bases were killed by a jet of steam applied near the base of the shoots for 3 to 6 minutes.) That the tissues were actually killed before the

P³²O₄ was applied was verified in all cases by the use of the resistance bridge described by DePlater and Greenham (1959). The tissues were considered to be dead when the high-frequency (1Mc/s):low-frequency (1kc/s) ratio approached 1.
Normal conditions of transport in the host occurred after P³²O₄ reached the host tissues. Extensive downward transport usually did not occur when the host branch was girdled directly above the

TABLE 6
EFFECT OF DEFOLIATION ON TRANSLOCATION OF C¹⁴-LABELED ASSIMILATES FROM TREATED WHITE FIR LEAVES TO LATERAL BRANCHES INFECTED BY DWARF MISTLETOE
(Radioactivity determined two weeks after labeling of terminal foliage of main branch)

Growth stage of host, and season	Intensity of autoradiographs*						
	Uninfected branch	Infected branch					
	Defoliated†	Not defoliated‡			Defoliated		
	Host stem	Host stem	Endophytic system	Mistletoe shoot	Host stem	Endophytic system	Mistletoe shoot
Dormant (fall).....	2	0	0	0	7	7	3
Dormant (winter).....	2	0	0	0	7	7	2
Dormant (spring).....	2	1	1	0	7	7	4
Early active (spring)....	7	7	7	7	7	7	7
Active (summer).....	7	0	0	0	7	7	7

* Radioactivity ratings are on a scale from 0 to 7, on the basis of counts per minute per mg. For a fuller explanation, see text, p. 118.
† Nondefoliated shoots usually 0, but others were noted and are described in the text.
‡ Other responses were noted and are described in the text, together with information on digger pine and lodgepole pine.

Fig. 8. Mounted plants (above) and autoradiographs (below) of white fir branches with lateral branches infected with *Arceuthobium campylopodum*. Terminal foliage (above arrows) was treated with $C^{14}O_2$, and collections were made two weeks later. A. infected branch not defoliated; treated August 31, 1960. B. infected branch defoliated just before treatment on August 31, 1960. Note that there was no translocation into basal section (X). C. infected branch not defoliated; treated November 20, 1960. Note upward translocation in first stem cross-section of infected branch (b).

point of treatment of the mistletoe shoots. The label evidently entered the xylem and was carried upward through the girdles but was kept from moving downward in the phloem (along with assimilates coming down from the leaves) because of the girdles. Limited downward movement below the girdle did occur, however, even though such movement was for a short distance and probably in the xylem.

Translocation of Labeled Herbicides, Phosphate, and Sulfate in Infected Branches

Infected white fir and digger pine branches were studied. Materials were applied to the host foliage, bark, and

wood, and to the aerial shoots of the dwarf mistletoes.

Only traces of 2,4-D acid migrated out of white fir leaves into the connecting branches; thus only traces, if any, reached the endophytic system or aerial shoots. All of the leaf applications shown in table 9 were on one-year-old leaves; however, a few tests on current-season white fir leaves (8/60) did not indicate any appreciable improvement in absorption and translocation. There was no indication that the ester of 2,4,5-T migrated appreciably out of the treated white fir leaves, even after nine months. Evidently 2,4,5-T is even more poorly absorbed and translocated than is 2,4-D. On the other hand, amitrole

Fig. 9. Mounted plants (above) and autoradiographs (below) of noninfected branches, terminal foliage of which (above arrows) was exposed to $C^{14}O_2$. Collections were made two weeks later. Lateral branches were defoliated to determine effect of this treatment on translocation. A. white fir with a lateral branch defoliated (d) just before treatment on July 25, 1961. B. digger pine with two lateral branches defoliated (d) and one branch covered with aluminum foil (a) just before treatment on May 11, 1962. Note abundant translocation into defoliated branches but very limited movement in covered branch. C. blue oak with two lateral branches defoliated (d) just before treatment on July 10, 1963.

was absorbed and translocated out of both young (not shown) and one-year-old leaves into the host stems and mistletoe. Both uptake and transport of amitrole were considerable when the treated host foliage was covered with a polyethylene bag (to produce a water-saturated atmosphere) for 24 hours immediately after treatment. Atrazine did not appear to be translocated downward, although some did move out of the treated leaves and was swept upward to the branch tips 6 to 10 cm away; this movement was likely in the xylem. Both $P^{32}O_4$ and $S^{35}O_4$ were absorbed and translocated downward in the stems, with considerable accumulation in the endophytic system and aerial shoots of

the dwarf mistletoes. Absorption and translocation were very greatly enhanced when the mature digger pine needles (current season) were scraped with a razor blade before the applications were made (fig. 12). It was also noted that branches with yellowish needles (which appeared to be dying) absorbed and translocated $P^{32}O_4$ considerably better than did branches with green, healthy needles.

Apparently only small amounts of labeled herbicides were absorbed through the intact bark. There was no evidence that the butoxy ethanol ester of 2,4,5-T applied in diesel oil was absorbed in a two-week period. However, some label clearly appeared in the bark

TABLE 7
EFFECT OF GIRDLING ON DOWNWARD TRANSLOCATION IN THREE
VARIOUSLY LABELED CONIFERS INFECTED BY DWARF MISTLETOE
(Radioactivity determined two weeks after labeling of foliage)

Conifer and label	Girdle			Translocation of label
	Type	Location	Date of girdling and labeling	
White fir ($C^{14}O_2$).....	Simple	Near base of stem	8/60	Into infection only
	Simple	Near base of stem	9/60	To girdle
	Simple	Near base of stem	5/61	Strong movement to girdle
	Simple	Near base of stem	7/61	No downward movement
	Simple	Near base of stem	8/62	Into infection only
	Double spiral	Above infection	7/62	To 8 cm below girdle
	Double spiral	Above infection	7/62	Trace to base of stem
	Loop	Above infection	8/62	Into infection only
	Fire	Above infection	8/62	To girdle
	Detached branch		8/60	Into infection only
	Stem base of mistletoe shoots		7/62	Only into unsteamed shoot
Lodgepole pine ($C^{14}O_2$)	Simple	Near base of stem	8/62	To 10 cm below girdle in pith
Digger pine: $C^{14}O_2$	Simple	Near base of stem	3/63	15 cm below girdle in resin canals
	Double spiral	Above infection	7/62	Through the infection
	Simple (2 trials)	Near base of stem	3/62	Slightly beyond girdle
	Simple	Side branch	12/61*	Down treated branch; through girdle on side branch
	Simple	Side branch	12/61	None up side branch

* Movement up ungirdled branches varied from none to strong on this date.

nine months after the applications were made. Amitrole was translocated downward only, and atrazine upward only, following summer applications to the bark. The quantity of label taken in was very small.

Application to the wood (through single cuts in the branches) was by far the most effective method of introducing labeled compounds for movement and accumulation in the same branches (table 9 and fig. 13); however, this method was not so effective with amitrole as were foliage applications when the leaves were covered with a polyethylene bag for 24 hours. Phosphate and sulfate were translocated upward following application to single cuts in the branches beneath the infections; such treatments resulted in a great accumulation of P^{32} and S^{35} by the endophytic systems. Basipetal translocation to the base of the branches occurred except when branches were girdled. Girdles

either directly above or below the point of application kept the label from being transported to the bases of the stems.

Two girdles, one directly above and the other directly below the infections, did not prevent a high degree of accumulation of label by the dwarf mistletoes (endophytic systems and aerial shoots), but did result in an extreme variation of the labeling within each endophytic system and between aerial shoots.

Labeling of the host stems and mistletoes was never completely uniform when the $P^{32}O_4$ and $S^{35}O_4$ were introduced into single cuts in the branches; however, it is significant that all parts of the host bark and dwarf mistletoes were labeled from such treatments (table 9).

No label entered the endophytic system or host tissues following application of C^{14} -labeled 2,4-D acid, amitrole, atrazine, and urea to dwarf mistletoe shoots on white fir, and of 2,4-D acid,

TABLE 8
TRANSLOCATION FOLLOWING TREATMENT OF DWARF MISTLETOE
SHOOTS WITH C¹⁴O₂ AND P³²O₄
(Radioactivity determined two weeks after labeling)

Conifer and label	Manipulation of mistletoe shoots	Date of labeling	Intensity of autoradiographs*			
			Host stem		Mistletoe	
			Above infection	Base of stem	Endophytic system	Untreated aerial shoots
White fir, digger pine, lodgepole pine (C ¹⁴ O ₂)	None (28 trials; all seasons)		0	0	0	0
White fir (P ³² O ₄).....	None (8 trials)	10/61	0	0	0	0
	None (2 trials)	10/61	1	0	7	0
Digger pine (P ³² O ₄).....	None (2 trials)	3/62 and 5/64	0	0	0	0
	Puncture (2 trials)	12/61	1 & 6	0 & 6	7 & 7	..
	Puncture; girdle below endoph. syst.	12/61	7	0	7	0
	Puncture; girdle above endoph. syst. (2 trials)	12/61	5 & 7	0 & 0	7 & 7	..
	Puncture; girdle above and below endoph. syst.	11/61	2	1	7	..
	Base steamed (3 trials)	3/62	0	0	0	0
	Base steamed	3/62	3	1	2	..
	Base steamed; punctured (2 trials)	3/63	0 & 3	0 & 1	6 & 7	..
	Puncture (5 reps.)	7/64	0	0	0	0
	None (4 reps.)	7/64	0	0	0	0
	None	7/64	0	0	6	..

* Radioactivity ratings are on a scale from 0 to 7, on the basis of counts per minute per mg. For a fuller explanation, see text, p. 118.

amitrole, MH, paraquat, 2,4,5-T amine, and urea to similar shoots on digger pine. Treatments were made directly following cessation of shoot growth by white fir and shortly before initiation of shoot growth by digger pine.

GREEN MISTLETOES

Basipetal Translocation of C¹⁴ in Host Branches

Representative distribution of C¹⁴ in green mistletoe-infected branches two weeks after the terminal foliage was exposed to C¹⁴O₂ is shown in table 10 and figure 14. Data are for four infected conifer and four infected dicotyledon branches and for some uninfected branches.

Downward translocation away from the treated region occurred at all seasons of the year. In only one instance did the C¹⁴ fail to reach the base of the stem within the two-week period of study. In the Macnab cypress in late

spring, downward movement ceased at 20 cm below the treated area, but label did accumulate in a cone directly beneath the treated area. Apical translocation was strong, as evidenced by the marked labeling of the male cones (microsporangiate strobili) at the ends of the branches.

The bark directly below the treated area was uniformly labeled around the stem, but the uniformity disappeared after side branches were encountered. Large branches were especially effective in restricting the C¹⁴ to a limited portion of the bark.

Small amounts of label (generally less than 1 cpm/mg) were usually

Fig. 10. Mounted plants (above) and autoradiographs (below) of white fir branches infected with *Arceuthobium campylopodum*; aerial shoots of mistletoe exposed to $C^{14}O_2$ on July 24, 1961. A, branch undisturbed; B, foliage of branch covered with aluminum foil during two-week period between treatment and harvest; C, branch defoliated at time of treatment. Note lack of basipetal translocation in A, B, and C.

present in the endophytic system. When this occurred, traces were also present in the aerial shoots.

Translocation of C^{14} Following Treatment of Mistletoe Shoots

Green mistletoe shoots were exposed to $C^{14}O_2$ in a manner similar to that used for host shoots. Translocation was comparable with that in green, nonparasitic plants (table 11 and fig. 15). Downward translocation from the treated foliage usually occurred, with the intensity of movement and accumulation varying considerably in different experiments. In some tests, the endophytic system was highly labeled two weeks after treatment, while in others it was only lightly labeled. In some tests there was no downward movement of label from

the treated part of the mistletoe shoots; however, upward movement to the shoot apices usually occurred as indicated by intense labeling of that area (fig. 15B).

Usually only a portion of a single green mistletoe shoot was exposed to the $C^{14}O_2$, and generally only parts of the endophytic system became labeled—or at least strongly so. Some of the assimilates in the less strongly labeled portions of the endophytic system and all of that in the unlabeled portions probably came from other mistletoe shoots (at least, for the two-week experimental period).

The untreated, lateral branches of the mistletoe shoots with terminal portions treated contained only traces of label, which evidently had moved into these lateral shoots from the main stem

Fig. 11. Mounted plants (above) and autoradiographs (below) of digger pine branches infected with *Arceuthobium campylopodum*; aerial shoots were treated with either $C^{14}O_2$ or $P^{32}O_4$, and collections were made two weeks later. A. mistletoe shoots pricked with pin in several places before $P^{32}O_4$ was applied November 24, 1961. B. mistletoe shoots (not pricked) treated with $P^{32}O_4$ on March 8, 1962. Note complete absence of any translocation into host. A portion of one of the less radioactive, treated shoots was mounted. C. mistletoe shoots treated with $C^{14}O_2$ on May 11, 1962. Note absence of any translocation into host.

of the treated mistletoe shoot in the process of moving toward the endophytic system. No label was present in green mistletoe shoots not connected with the treated mistletoe shoot.

Little label appeared to move from the green mistletoes into the host tissues, and any that did was in the immediate vicinity of the endophytic system.

Translocation of C^{14} in Infected, Defoliated Lateral Branches

Branches infected by green mistletoe often die above the infections. Such branches are often completely free of host foliage and must obtain their nutrition either from the mistletoes or from other host branches. Such branches were

studied in order to determine the source of their nutrition. A sufficient number of branches in this condition were not readily accessible, and were therefore prepared by pruning. Blue oak stems were prepared one year in advance of the studies, and Macnab cypress branches four months in advance.

Autoradiographs indicate that most of the assimilates required by the defoliated branches came from other host branches and that nearly all of the assimilates required by the green mistletoes was produced by them (table 12 and fig. 16). Traces of assimilates moved from the defoliated branches into the endophytic system and aerial shoots of the mistletoe; this label was present in

TABLE 9
AVERAGE TRANSLOCATION OF VARIOUS C¹⁴-LABELED HERBICIDES,
PHOSPHATE, AND SULFATE, FOLLOWING DUPLICATE AND TRIPPLICATE
APPLICATIONS TO LEAVES, BARK, AND WOOD OF WHITE FIR AND
DIGGER PINE INFECTED WITH DWARF MISTLETOE
(Radioactivity determined two weeks after labeling unless otherwise noted)

Labeled material	Plant species	Date of labeling	Intensity of autoradiographs*			
			Host stem		Mistletoe	
			Above infection	Base of stem	Endophytic system	Aerial shoots
Leaf application						
2,4-D acid.....	white fir	8/60	1	0	1	0
2,4,5-T ester.....	white fir	8/61	0	0	0	0
2,4,5-T ester.....	white fir (9 months)	8/61	0	0	0	0
Amitrole.....	white fir	8/60	1	0	1	1
Amitrole.....	white fir†	8/60	3	1	3	2
Atrazine.....	white fir	8/60	1	0	0	0
Phosphate.....	digger pine	12/61	3-7	2-4	7	4-7
Phosphate.....	digger pine	3/62	2	1	2	1
Phosphate.....	digger pine (5 reps.)	6/64	4	2	4	1-4
Phosphate.....	digger pine (leaves scraped before 5 reps.)	6/64	7	6	7	4-7
Bark application (below infections)						
2,4-D acid.....	white fir	8/60	0	1	1	0
2,4,5-T ester.....	white fir	8/61	0	0	0	0
2,4,5-T ester.....	white fir (9 months)	8/61	1	0	1	1
Amitrole.....	white fir	8/60	0	1	0	0
Atrazine.....	white fir	8/60	1	0	1	0
Wood application (to cut below infections)						
2,4-D acid.....	white fir	8/60	1-3	1-2	1-3	1-2
Amitrole.....	white fir	8/60	1	0	1	0
Atrazine.....	white fir	8/60	0-3	0	0-1	0
Phosphate.....	digger pine (girdle below endophytic system)	3/62	7	0	7	2-7
Phosphate.....	digger pine (girdles above and below endophytic system)	3/62	1-7	0	0-7	0-7
Phosphate.....	digger pine	3/62	7	7	7	7
Sulfate.....	digger pine	3/62	4-7	0-4	4-7	2-7
Sulfate.....	digger pine (girdle below endophytic system)	3/62	5-7	0	5-7	5-7
Sulfate.....	digger pine (girdles above and below endophytic system)	3/62	5-7	0	5-7	5-7

* Radioactivity ratings are on a scale from 0 to 7, on the basis of counts per minute per mg. For a fuller explanation, see text, p. 118.
† Treated area covered with polyethylene bag for 24 hours.

low concentration in the outer portion of the stems of the mistletoe shoots. When the mistletoe shoots were exposed to C¹⁴O₂, a considerable amount of labeled assimilates migrated to the endophytic system (fig. 16), which showed dark on autoradiographs. Traces of label may possibly have passed from the endophytic system into the adjacent host tissues, but this was not demonstrated with certainty. Girdling the defoliated branches to deprive them of any opportunity to obtain food from other branches still did not enable them

to obtain assimilates from the mistletoes.

Defoliation of uninfected branches of Macnab cypress resulted in very little of the assimilates flowing into them during the dormant season, but a considerable amount did flow into them during the growing season. These results were similar to those obtained on the mistletoe-infected branches. In all tests, defoliation of uninfected blue oak branches resulted in appreciable movement of assimilates to them.

Translocation of Labeled Phosphate Following Application to Host Foliage, Cuts in Host Stem, and Mistletoes

Studies were conducted on the absorption and translocation of $P^{32}O_4$ because this chemical is as freely mobile in plants as any herbicide is likely to be. Furthermore, it must be applied in a manner similar to that used with herbicides.

Some typical results from these studies are presented in table 13 and figure 17. Applications to the foliage of Macnab cypress and blue oak resulted in small amounts of label entering the mistletoe shoots. The intensity of such labeling was slightly greater than that obtained with assimilates (table 12). It was found that $P^{32}O_4$ migrated up lateral branches in response to defoliation in much the same manner as did assimilates (table 12).

The most effective method of introducing label into the green mistletoe was to apply it to the host stem below the infections (table 13). These treatments resulted in not only the host stems but also the mistletoe shoots becoming very strongly labeled. During the winter period, the green mistletoe shoots on Macnab cypress were appreciably stronger-labeled than was the host foliage.

Although the endophytic system was usually strongly labeled following ap-

Fig. 12. Mounted plants (above) and autoradiographs (below) of digger pine branches infected with dwarf mistletoe. Foliage was treated with $P^{32}O_4$ on July 1, 1964, and collections were made one week later. Treated foliage (above arrows) was too radioactive to use, and was discarded to prevent contamination. A through E: leaves were severely scraped with a razor blade before $P^{32}O_4$ was applied. Note abundant translocation in all five branches. F through J: the $P^{32}O_4$ was applied to leaves without scraping.

plication of $P^{32}O_4$ to the mistletoe foliage (table 13), labeling was stronger when the applications were made to cuts in the stems. Furthermore, in numerous instances (mid-spring) only small amounts of label appeared to migrate to the endophytic system (fig. 17). In one instance, considerable label entered the host stem following application to green mistletoe on Macnab cypress.

Fig. 13. Mounted plants (above) and autoradiographs (below) of digger pine branches infected with *Arceuthobium campylopodum*. $P^{32}O_4$ and $S^{35}O_4$ were applied to cuts in stems below infections at points indicated by arrows. Treatment period was two weeks. A. $P^{32}O_4$ applied to a single cut in stem on January 3, 1962, beneath girdle (g). Except for section directly beneath point of application, no label was present in stem cross-section below that point. Basal section is indicated by (x). B. $S^{35}O_4$ applied to a single cut in stem on March 23, 1962. No girdle was used. Label was present in stem cross-sections beneath point of treatment.

TABLE 10
SELECTED EXAMPLES OF DISTRIBUTION OF C¹⁴-LABELED ASSIMILATES IN
NORMAL BRANCHES OF VARIOUS TREE SPECIES, AND IN
BRANCHES INFECTED WITH GREEN MISTLETOE
(Radioactivity determined two weeks after labeling of terminal foliage)

Plant species and stage of growth	Date of labeling	Intensity of autoradiographs*			
		Host stem		Mistletoe	
		Above infection	Base of stem	Endophytic system	Aerial shoots
Blue oak:					
Active.....	5/63	7	2-7	1	1
Active†.....	5/63	7	2-6	1	..
Active.....	5/64	7	7	2	0
Postactive.....	7/61	7	1-7	1	1
Postactive.....	7/63	7	1-7	..	..
Postactive‡.....	8/63	7	2-7	1	..
California black oak:					
Postactive.....	8/61	6	6	1	1
Interior liveoak:					
Dormant.....	3/62	7	2-7	1	1
California black walnut:					
Active.....	6/61	7	4-7	1	1
Incense cedar:					
Active.....	8/62	6	0-5	0	0
Western juniper:					
Active.....	7/61	7	0-7	0	0
White fir:					
Active.....	7/61	7	3-7	1	1
Macnab cypress:					
Dormant.....	3/62	7	1-5	1	1
Active§.....	5/62	7	0	..	..
Active.....	5/62	7	0-5	..	..
Active.....	5/64	7	7	2	2
Active.....	7/62	7	7	0	0
Postactive.....	8/62	5-7	3-5	1	0

* Radioactivity ratings are on a scale from 0 to 7, on the basis of counts per minute per mg. For a fuller explanation, see text, p. 118.

† Mistletoe shoots removed; endophytic system covered with aluminum foil.

‡ Mistletoe shoots removed.

§ Label into cone below treated leaves.

Translocation of 2,4,5-T Ester and Paraquat Applied to Mistletoes

Paraquat applied to mistletoe on Macnab cypress in May did not translocate detectably within the shoots. The butoxy ethanol ester of labeled 2,4,5-T applied to mistletoe on blue oak translocated detectably for only 3 to 4 cm. Nonradioactive mixtures of 2,4-D and 2,4,5-T did not translocate in herbicidal proportions except over very short distances. When these mixtures were applied to the base of mistletoe shoots (six replications) the infections were killed in two instances and the mistletoes plus the host branches were killed in four

instances. When the applications were made to the terminal part of the mistletoe shoots, only a limited portion of the aerial shoots was killed.

Photosynthesis and Respiration

Photosynthesis has been shown to occur in the dwarf mistletoe (Hull and Leonard, 1964b), but at a rate insufficient to be detectable by the Beckman infrared gas analyzer. It was detectable, however, by means of C¹⁴O₂.

Limited studies were conducted on the photosynthetic capacity of the green mistletoe (table 14). The mistletoe shoots were removed from host trees,

Fig. 14. Mounted plants (above) and autoradiographs (below) of branches infected with green mistletoe (*Phoradendron*). Terminal foliage was treated with $C^{14}O_2$ and collections were made two weeks later. Note absence of label in endophytic system (e). A. California black walnut infected with *P. flavescens* var. *macrophyllum*, treated June 6, 1961. B. western juniper infected with *P. juniperinum* var. *ligatum*, treated August 8, 1961. C. blue oak infected with *P. flavescens* var. *villosum*, treated August 29, 1961.

and portions of the shoots were placed with their bases in water. They were immediately transferred to a chamber in which the atmosphere was monitored by an infrared gas analyzer, for photosynthesis and respiration studies. Green mistletoe apparently requires about 800

foot-candles of light for photosynthesis to balance respiration. Greenhouse-grown blue oak and interior liveoak required less light to achieve such balance than did green mistletoe; the oaks also fixed more carbon per unit of fresh weight than did the mistletoe.

COMPARISON OF TRANSLOCATION DATA: DWARF AND GREEN MISTLETOES

Normal and Mistletoe-infected Host Trees

There was no indication that the green mistletoes influenced translocation of C^{14} in infected branches. This was also true with dwarf mistletoe except during the summer. Most of the host shoot

growth comes from food supplied by old needles—reserves and current assimilates (in red pine, Kozlowski and Winget, 1964). The additional drain made on the food supplied by dwarf mistletoe to nourish its endophytic system, aerial shoots, flowers, and fruit can result in no food available for trans-

TABLE 11
SELECTED EXAMPLES OF TRANSLOCATION OF C¹⁴-LABELED ASSIMILATES
IN GREEN MISTLETOE SHOOTS TWO WEEKS FOLLOWING LABELING

Host and stage of growth	Date of labeling	Intensity of autoradiographs*			
		Host stem		Mistletoe	
		Above infection	Base of stem	Endophytic system	Aerial shoots†
Blue oak:					
Dormant.....	2/63	0	0	0	0
Dormant‡.....	2/63	0	0	1	0
Dormant§.....	2/63	0	0	1	0
Active.....	5/63	0	0	5-7	0
Active.....	5/64	0	0	7	1
Postactive.....	7/61	0	0	0-7	0
Postactive.....	8/63	0	0	0-7	0
California black oak:					
Dormant.....	3/62	0	0	2-4	0
Interior liveoak:					
Dormant.....	3/62	0	0	2-4	0
California black walnut:					
Bud-break.....	3/61	0	0	7	0
Early growth.....	4/61	0	0	1-7	0
Full leaf.....	6/61	0	0	0	0
Incense cedar:					
Active.....	8/62	0	0	5	0
White fir:					
Active.....	8/62	0	0	0-7	1
Macnab cypress:					
Dormant.....	3/62	0	0	0-6	0
Active.....	5/64	0	0	7	1
Active.....	7/62	0	0	0-7	0
Postactive.....	8/62	0	0	0-7	0

* Radioactivity ratings are on a scale from 0 to 7, on the basis of counts per minute per mg. For a fuller explanation, see text, p. 118.
† Treated portion of green mistletoe shoots was always 7 (dark autoradiograph). Data are for shoots not connected with treated shoots.
‡ Branch collected 44 days after labeling of mistletoe.
§ Branch collected 80 days after labeling of mistletoe.

port basally beyond the infected areas during the summer.

The downward rate of translocation in white fir branches (whether infected by dwarf mistletoe or not) was considerably faster during the winter than during the summer and fall (table 5). The rapid rate continued until after the beginning of shoot growth in the spring. This marked drop also occurred in infected red fir, ponderosa pine, and Jeffrey pine (table 3). Nelson (1964) reported translocation to the roots of conifers before growth began, but not when shoot growth was rapid.

The slow rate of translocation in white fir in the fall may have been the result of moisture stress on photosynthesis (Kramer, 1964; Clark, 1961) or directly upon translocation (Kramer, 1964). Summer rainfall is low in the Sierra Nevada, and could cause moisture stresses in the trees.

The evidence indicates that dwarf mistletoe infections do not act as physical blocks to translocation. When such blocks do occur, they appear to be temporary and to depend on the capacity of the mistletoes to compete with the host tissues for food. Thus such blocks are strong sinks. The translocation of assimilates presumably will be most frequently blocked by (1) large, active infections, (2) unusually large requirements by host tissues because of vigorous growth, and (3) reduced photosynthetic activity of the host, whether temporary or permanent.

Fig. 15. Mounted plants (above) and autoradiographs (below) of branches infected with green mistletoe (*Phoradendron*). Terminal portion of mistletoes (above arrow) was exposed to $C^{14}O_2$; collections were made two weeks later. A. *P. flavescens* var. *macrophyllum* on California black walnut treated March 17, 1961, at time of bud-break of the walnut. Note translocation into endophytic system of mistletoe. B. same as (A) above, but treated June 6, 1961. Note absence of any downward translocation in mistletoe shoots. C. *P. juniperinum* var. *libocedri* on incense cedar treated August 17, 1962. Note downward transport into endophytic system.

Translocation into Dwarf and Green Mistletoes

Green mistletoes accumulated no assimilates (or only traces) from their hosts. When traces of label did appear, they may have arrived via the xylem. Miller (1938) cited several references on the occurrence of sugar in the xylem sap of trees, especially in the spring.

In contrast, the dwarf mistletoes obtained large quantities of photosynthate from their hosts at all seasons of the year. The storage of food reserves in the endophytic system during the winter enables dwarf mistletoes to grow rapidly when conditions become suitable. The endophytic system probably

grows very little on conifers in the high Sierra Nevada during the winter. However, the endophytic system in digger pine (present at low elevations; see table 1) grows about as rapidly during winter as during summer (Scharpf, 1962b). The accumulation of food reserves by the endophytic system during winter was about the same on digger pine (low elevation) as on white fir (high elevation). This similarity in food storage suggests that growth of the endophytic system in white fir does not stop completely during winter.

Most of the food materials are probably absorbed by the endophytic system of dwarf mistletoes at the point where

TABLE 12
SELECTED SAMPLES OF TRANSLOCATION OF C¹⁴-LABELED ASSIMILATES
BETWEEN GREEN MISTLETOES AND THEIR HOSTS AS INFLUENCED
BY DEFOLIATION OF INFECTED BRANCHES
(Radioactivity determined two weeks after labeling)

Growth stage and treatment of host and mistletoe	Intensity of autoradiographs*					
	Defoliated			Not defoliated		
	Host stem	Mistletoe		Host stem	Mistletoe	
		Endophytic system	Aerial shoots		Endophytic system	Aerial shoots
Macnab cypress:						
Host foliage treated†						
Dormant.....	2	1	1	1	1	1
Active.....	7	2	1	0	0	0
Active (base host stem girdled)‡ ..	7	2	1	..	..	..
Mistletoe treated						
Dormant.....	0	4	0§	0	4	0
Active.....	0	7	1	0	7	1
Active (base host stem girdled) ...	0	7	1	..	..	..
Blue oak:						
Host foliage treated†						
Active (3 reps.).....	7	1	0	0	0	0
Mistletoe treated						
Active (3 reps.).....	0	7	1‡	0	0	0

* Radioactivity ratings are on a scale from 0 to 7, on the basis of counts per minute per mg. For a fuller explanation, see text, p. 118.
† Host foliage on uninfected main stem. Data on autoradiography are for infected lateral branches.
‡ Girdle was below point at which defoliated infected branch joined normal branch treated with C¹⁴O₂.
§ Nontreated lateral shoots.
|| Girdle at base of infected branch.

the filamentous strands and sinkers pass through the active phloem (fig. 18A). Other details on anatomy may be found in Kuijt (1960), Srivastava and Esau (1961), and Scharpf (1963a).

The aerial shoots of dwarf mistletoes accumulated label most actively during the summer, possibly because of the requirements for shoot growth and flower production on both male and female plants, and for fruit development on the females. High levels of label accumulated in the fruit.

Removal of the dwarf mistletoe shoots resulted in enhanced accumulation of label by the endophytic system (table 4 and Hull and Leonard, 1964b), whereas covering the shoots with aluminum foil had little effect (table 4). The increased accumulation may have been due to removal of a sink or it may have resulted from a shift in the hormone balance following shoot removal.

Removing the aerial shoots of green mistletoes did not induce the endophytic system to accumulate more than traces of label from the hosts (table 10). Whether these traces have any nutritional significance remains to be determined. In this connection, Wagener (1925) reported that a green mistletoe on incense cedar had apparently remained alive in the bole for 100 years, without aerial shoots.

Translocation within and between Mistletoes and Hosts

Dwarf and green mistletoes both carry on photosynthesis, but the former does so at a rate far below that necessary for its own nutrition (Hull and Leonard, 1964b). In green mistletoes, photosynthesis is in excess of respiration, but the CO₂-compensation point appears to be greater than that of the host foliage (table 14). This relation-

Fig. 16. Mounted plants (above) and autoradiographs (below) of blue oak branches infected with green mistletoe (*Phoradendron*). All shoots with foliage on infected branches were removed before treatment. Host foliage (above arrows on nondefoliated branch) and mistletoe shoots (above arrow on defoliated branch) were exposed to $C^{14}O_2$; collections were made two weeks later. A. host foliage treated May 25, 1964. Note translocation up infected lateral branch (d), but near-absence of label in mistletoe. Infected branch was defoliated 11 months before experiment. B. control. Note faint pseudoautoradiographs. C. same as (A), with appreciable label in a new host branch (b) that developed during two-week treatment period. D. host foliage treated July 1, 1963, and collected 10 days later. Defoliated branch (d) had mistletoe shoots and leaf-containing host shoots removed 33 days before treatment. E. mistletoe shoot (above arrow) treated May 24, 1964, on a branch free of host foliage. Branch defoliated 11 months before treatment with $C^{14}O_2$. F. same as (E), except girdled (g) as well as defoliated.

ship is suggested by the fact that the largest green mistletoe plants are located on their hosts in places where they receive appreciable light—for example, those on the tops of white fir (fig. 19). Apparently no such relationship occurs on deciduous hosts, probably because the green mistletoes receive several months of full sunlight during the leaf-free period or dormant season. On incense cedar and Macnab cypress the relationship also fails to hold, but the host foliage is quite open and light intensities are high throughout the canopies.

Dwarf mistletoes receive almost all of

their organic carbon from their hosts, whereas green mistletoes apparently receive only traces. In dwarf mistletoe the carbon is picked up by the endophytic system, in which it may become quite concentrated. It moves from the endophytic system into the aerial shoots at all times of the year, the rate of movement being least in winter and greatest during late spring and summer. The path of translocation must be in living tissues, since steaming the bases of the shoots prevented any movement of assimilates into them. The label concentrates at the nodes of the aerial shoots at all times of the year, and is especially

TABLE 13

SELECTED EXAMPLES OF TRANSLOCATION OF $P^{32}O_4$ FOLLOWING
APPLICATION TO FOLIAGE AND CUTS IN STEMS OF HOST PLANTS,
AND TO GREEN MISTLETOES

(Treatment was for seven days during period of active growth)

Point of application	Intensity of autoradiographs*			
	Host stem		Mistletoe	
	Above infection	Below infection	Endophytic system and host stem†	Aerial shoots
Macnab cypress:				
Foliage.....	7	2	7	2
Cut in stem below endophytic system.....	7	6	7	7
Mistletoe.....	1	1	7	..
Blue oak:				
Foliage.....	7	7	7	2
Cut in stem below endophytic system.....	7	7	7	7
Mistletoe.....	0	0	2	..

* Radioactivity ratings are on a scale from 0 to 7, on the basis of counts per minute per mg. For a fuller explanation, see text, p. 118.

† Reading includes both host and mistletoe tissues since no clear separation can be made when using P^{32} .

concentrated in the male and female flowers and in the fruit. Cross-sections of the shoots show a uniform concentration of label except at the center, where xylem is located and concentration is less intense (fig. 18B).

Exposure of both dwarf and green mistletoe shoots to $C^{14}O_2$ in sunlight resulted in the formation of labeled sugars. From 40 to 50 per cent of the label in dwarf mistletoe shoots was found to be sucrose (Hull and Leonard, 1946b). No study of sugar content was made on green mistletoes.

Numerous experiments conducted at all times of the year failed to show any indication of basipetal flow of labeled assimilates in dwarf mistletoe shoots. Furthermore, such flow could not be induced by covering the host foliage with aluminum foil or by defoliation to create carbohydrate deficiencies in the host (fig. 10). Upward translocation occurred in the exposed shoots, and resulted in the same distribution of label as that translocated through the endophytic system. Label was present in the outermost parts of the shoots, however,

in contrast to the more general distribution when label came from the endophytic system.

In green mistletoes, translocation was generally downward, with movement into the endophytic system (fig. 15 and table 11). In many instances the apices, flowers, and fruit in the $C^{14}O_2$ -treated part of the shoot were more intensely labeled than were other parts of the treated area. This type of labeling suggests that both upward and downward movement had occurred in the same shoots. In one experiment the movement was entirely apical (fig. 15B). Downward transport resulted in varying degrees of labeling of the endophytic system in different experiments (table 11), probably reflecting the relative competition for food by the mistletoe shoots and the endophytic systems. Usually only a portion of the endophytic system was labeled by treatment of a single shoot. This was to be expected since other shoots could nourish those parts of the endophytic system with which they were most closely associated.

Movement of assimilates from the

Fig. 17. Mounted plants (above) and autoradiographs (below) of blue oak branches infected with *Phoradendron flavescens* var. *villosum*. Autoradiographs indicate translocation of $P^{32}O_4$ during a two-week period following its application to foliage and cuts in the stem of the host, and to mistletoe shoots. Applications were made May 25, 1964. A. host foliage treated (above arrow—discarded). Note limited uptake by endophytic system (e). B. mistletoe shoot treated (above arrow—discarded). Note absence of any movement of label into host from mistletoe. C. application made to a cut in stem (arrow). Note intense movement into both host and parasite above point of application.

green mistletoe into the host branches was not significant, but traces of label were present in tissue directly associated with the endophytic system. No evidence was obtained that assimilates could move from dwarf mistletoes into their hosts. Evidently, materials within both dwarf and green mistletoes tend to remain there.

Nutrition of Foliage-free, Infected Branches

The current studies have thrown much light on the effect of mistletoes on the nutrition of infected, defoliated branches. Mistletoe-infected, defoliated host branches received nourishment exclusively from normal host branches. Such branches infected with dwarf mistletoes imported assimilates at all seasons of the year; those infected with green mistletoes imported appreciable assimilates (or $P^{32}O_4$) only during the growing season. The behavior of branches infected with green mistletoe was identical with that of similar but noninfected branches with respect to the importation of labeled assimilates or $P^{32}O_4$. Apparently neither the green nor the dwarf mistletoes contributed to the nutrition of host branches that were free of foliage.

The nutrition of defoliated infected and noninfected branches is similar to that of stumps connected to trees by root grafts, which may remain alive for years and continue to grow. Bormann and Graham (1959) describe such grafting in eastern white pine.

TRANSLOCATION AND POTENTIAL CHEMICAL CONTROL

For successful chemical control of mistletoes, herbicides must be toxic to the parasites and noninjurious to hosts. Such herbicides should be readily absorbed, translocated, and slowly degraded. Differences in the organic nu-

trition of dwarf and green mistletoes might determine how such sprays could best be applied.

Foliar application is desirable because an area can be treated rapidly with moderate uniformity and at a relatively

TABLE 14

CHANGES IN CO₂ CONCENTRATION OF AIR IN 10 MINUTES BY 10 GRAMS
OF DETACHED SHOOTS OF GREEN MISTLETOE, BLUE OAK,
AND LIVEOAK UNDER DIFFERENT LIGHT INTENSITIES

Plant species	Changes in CO ₂ concentration at:		
	0 ft.-c.	800 ft.-c.	10,000 ft.-c.
	<i>per cent</i>	<i>per cent</i>	<i>per cent</i>
Green mistletoe.....	+5	0	-7
Blue oak.....	+5	-8	-28
Interior liveoak.....	+8	-10	-33

low cost. Ideally, with dwarf mistletoe infections, the therapeutants should be applied when the aerial parts of the hosts are dormant, so that a greater percentage of the material will accumulate in the mistletoes. Winter and spring are preferable application seasons because translocation rates in conifers are faster (table 5, white fir) at those times, and relative humidity is likely to be higher. (High relative humidity favors absorption.)

Foliar applications on hosts infected with green mistletoe would be best during winter for deciduous hosts, and early in the growing season for evergreen hosts. The determining factor on deciduous hosts is that the absence of host foliage would permit coverage of the mistletoes by the sprays. With evergreen hosts, spray applications must be made at a time that is best for absorption and translocation in the mistletoes—apparently during rapid shoot growth of the hosts. Foliar applications would be most effective on infections near the ends of branches and at the tops of trees.

Direct application of sprays to the bark presents at least two problems: (1) the living tissues in the bark must be able to tolerate high concentrations of the therapeutants at the point of application; (2) only a low percentage of the material is able to penetrate bark, except where it is thin. Quick (1964) found that the iso-octyl ester of 2,4,5-TB was promising for controlling dwarf mistletoes only when the infections themselves were sprayed.

Therapeutants may be applied directly to host wood as a control for both dwarf and green mistletoes. This method is being used in Australia to control mistletoe on eucalyptus (Brown, 1959). In our research it was found to be a highly effective method of introducing P³²O₄ and S³⁵O₄ into dwarf and green mistletoes. Evidence from several sources indicates that in California the optimum period for application might extend from winter dormancy through active shoot growth (Leonard and Harvey, 1965). Such applications should be made to relatively moist wood at depths sufficient to avoid injury to the host bark. Possible injury should be further reduced by increasing the number of points of application per stem (Greenham and Brown, 1957). The current research showed a marked difference in concentration of the label in different parts of the host branches and mistletoes following single-point applications of P³²O₄ and S³⁵O₄ to the host wood. Single-point applications of P³²O₄ to 20 cm-diameter blue oak resulted in 75 to 100 per cent of the foliage becoming labeled in two weeks (Leonard, Glenn, and Bayer, 1965); however, at least a 500-fold difference in concentration of the label was found in different leaves. Winter application of therapeutants to mistletoe-infected evergreen trees should be less injurious than applications during the growing season. Less injury should result within a given time period because of the lower temperatures and metabolism rates, which per-

Fig. 18. Photomicrographs of dwarf mistletoe. A. cambial region of red fir infected by *Arceuthobium campylopodum*. Arrow points to a sinker passing through host phloem. It is here that most of the food exchange between host and mistletoe occurs. B. cross-section of aerial shoot of *A. americanum* on lodgepole pine. Arrow points toward central cylinder, central portion of which is xylem.

mit time for redistribution of the therapeutants before injury can develop. Greenham and Brown (1957) noted that a considerable redistribution of $F^{59}Cl_3$ occurred in the course of one year in mistletoe-infected eucalyptus. In view of the above considerations, multipoint deep application of therapeutants to the wood of mistletoe-infected evergreens during the winter shows promise because it should (1) result in minimum damage to the bark at the point of application; (2) reduce the concentration of the therapeutants received by leaves and branches; (3) allow time for redistribution when host tissues are least likely to be damaged; and (4) present no hazard to adjacent vegetation. A further advantage is that very large trees could be safely and easily treated. The same method should be feasible for deciduous trees, but with applications following the rapid spring growth, to reduce possible damage to developing tissues.

Single-point applications are especially attractive as a potential control method from the economic point of view. A single injection per stem of a toxic herbicide, picloram, in winter, was adequate to control madrone (*Arbutus menziesii*) and to kill most of the foliage on tan oak (*Lithocarpus densiflora*) but not on interior liveoak (Leonard and Hull, unpublished). Vité and Rudinsky

(1959) indicate that marked variation appears in the translocation pattern of acid fuchsin applied to the bases of trees, with the complete circumference being colored in some species. However, acid fuchsin is not subject to the same redistribution that is possible with highly phloem-mobile substances such as $P^{32}O_4$, $S^{35}O_4$, and maleic hydrazide. It is suggested that the potential for redistribution of highly mobile, low host-toxic therapeutants is far greater than the dye experiments of Vité and Rudinsky (1959) indicate. Single-point application is sufficiently promising to be worth considerable exploration.

The ideal therapeutant will probably never be found. Host tissues are usually not completely immune to selective materials, and their possible toxicity to the hosts must be considered. Applications to the wood and to dormant foliage should be more selective than those made to foliage during the growing season, or to the bark. Substances entering the dwarf and green mistletoes remain within them, but the parasites do not represent an important avenue of detoxification of therapeutant with respect to the host. The main hope of reducing toxicity to the host lies in obtaining sufficient redistribution of the therapeutant before localized injury can become important.

SUMMARY

Eight hosts of dwarf mistletoe (*Arceuthobium* spp.) and eight hosts of green mistletoe (*Phoradendron* spp.) were studied at different times of the year, following application of various labeled substances to the host foliage, bark, and wood, and to the mistletoe shoots. Radioactive materials used were $C^{14}O_2$, $P^{32}O_4$, $S^{35}O_4$, and C^{14} -labeled herbicides.

Dwarf mistletoes were found to derive their organic nutrition largely from their hosts, whereas green mistletoes manufactured their own carbohydrates.

The endophytic system and aerial shoots of dwarf mistletoe accumulated labeled assimilates from hosts at all seasons of the year. Under the same conditions, only traces of label were found in green mistletoes.

Dwarf mistletoe shoots carried on limited photosynthesis, but no labeled assimilates migrated basally in the endophytic system. In green mistletoe, assimilates did migrate basally to the endophytic system, but not to the hosts.

In treatment of dwarf mistletoe shoots with $P^{32}O_4$ results were similar, but

Fig. 19. Trees infected with green mistletoe (*Phoradendron*). A. *P. bolleanum* var. *pauciflorum* on white fir. Note dead top, heavy mistletoe infection (arrow). B. *P. flavescens* var. *villosum* on California white oak. Note random distribution of large infections over deciduous oak. C. *P. juniperinum* var. *libocedri* on incense cedar. Large infections occur randomly over this relatively open-crowned evergreen. D. *P. flavescens* var. *villosum* on California black oak. Large infections occur randomly over this deciduous oak.

some migration from shoots to hosts occurred—evidently via the apoplast, since steaming the shoot bases did not prevent such movement.

Defoliated host branches infected by either dwarf or green mistletoe obtained food supplied from other host branches, not from the mistletoes. During winter and spring, defoliated branches with dwarf mistletoe imported large quantities of label, but those with green mistletoe imported very little.

Except in the period shortly after bud-break in the spring, translocation in the hosts was not appreciably affected by either type of mistletoe infection. At that time, however, label migrated (in two weeks) from treated foliage down stems into, but not beyond, the mistletoe infection.

Only traces of label were absorbed and translocated into dwarf mistletoe following application of labeled herbicides to host bark.

Poor absorption and translocation occurred with labeled herbicides applied to white fir foliage. The only appreciable movement from treated foliage to dwarf mistletoe occurred with amitrole, and then only under high humidity. Rates of translocation of assimilates were similar in infected and noninfected branches of white fir. Rates were highest in winter and spring, lowest in summer and fall.

A light intensity of about 800 foot-candles was required to balance respiration and photosynthesis in green mistletoe.

ACKNOWLEDGMENTS

The research was supported by Grant 13070 from the National Science Foundation.

We wish to thank the U.S. Forest Service for making available the facilities of the Stanislaus Branch Station at Strawberry. We should also like to thank the California Division of Beaches and Parks for permitting study of dwarf mistletoe-infected digger pine in Mt. Diablo State Park.

Our thanks are also due Mr. C. G. Greenham, Plant Industry Division,

CSIRO, Canberra, Australia, for his role in some aspects of the current study; Mr. H. R. Offord, for his support of this project from its initiation, and Mr. Clarence R. Quick, both of the Pacific Southwest Forest and Range Experiment Station, Berkeley. Dr. J. R. Parmeter, Jr., University of California, Berkeley, supplied much useful background information concerning the mistletoes, and Drs. H. J. Rediske and K. R. Shea, of the Weyerhaeuser Company, made helpful suggestions.

LITERATURE CITED

ANONYMOUS

1962. White pine blister rust control. Ann. Rept. U.S.D.A. Forest Service, Region One. 1380 (3000). Missoula, Montana. 57 pp.
- BORMANN, F. H., and BEN F. GRAHAM, JR.
1959. The occurrence of natural root grafting in eastern white pine, *Pinus strobus* L., and its ecological implications. Ecology 40 (4): 677-91.
- BROWN, A. G.
1959. Mistletoe control on a large scale. Jour. Australian Inst. Agr. Sci. 25 (4): 282-86.
- CLARK, JOHN
1961. Photosynthesis and respiration in white spruce and balsam fir. State Univ. College of Forestry Tech. Bul. 84. Syracuse, N.Y. 72 pp.
- DEPLATER, C. V., and C. G. GREENHAM
1959. A wide-range A. C. bridge for determining injury and death. Plant Physiol. 34 (6): 661-67.
- GILL, L. S., and F. G. HAWKSWORTH
1961. The mistletoes. A literature review. U.S.D.A. Tech. Bul. 1242. 87 pp.
- GRASER, H. I.
1954. Methods for control of mistletoe; spot spraying in winter proves effective. Diamond Walnut News 36: 10.
- GREENHAM, C. G.
1964. Known and potential hazards to forest production by the mistletoes and dwarf mistletoes. Joint FAO-International Union of Forest Research Organizations. Rome. (Mimeo.)
- GREENHAM, C. G., and A. G. BROWN
1957. The control of mistletoe by trunk injection. Jour. Australian Inst. Agr. Sci. 23 (4): 308-18.
- GREENHAM, C. G., and O. A. LEONARD
1964. The amino acids of some mistletoes and their hosts. Amer. Jour. Bot. 52 (1): 41-47.
- HARTIGAN, D. T.
1949. Control of mistletoe. Australian Jour. Sci. 11: 174.
- HULL, RICHARD J., and OLIVER A. LEONARD
1964a. Physiological aspects of parasitism in mistletoes (*Arceuthobium* and *Phoradendron*). I. The carbohydrate nutrition of mistletoe. Plant Physiol. 39 (6): 996-1007.
1964b. Physiological aspects of parasitism in mistletoes (*Arceuthobium* and *Phoradendron*). II. The photosynthetic capacity of mistletoe. Plant Physiol. 39 (6): 1008-1017.
- KADAMBI, K.
1954. On Loranthus control. Indian Forester 80 (8): 493-95.
- KOZLOWSKI, THEODORE T., and CARL H. WINGET
1964. The role of reserves in leaves, branches, stems, and roots on shoot growth of red pine. Amer. Jour. Bot. 51 (5): 522-29.
- KRAMER, PAUL J.
1964. The role of water in wood formation. In: The formation of wood in forest trees, Martin H. Zimmermann, ed. New York and London: Academic Press. 519-32.
- KUIJT, JOB
1960. Morphological aspects of parasitism in the dwarf mistletoes (*Arceuthobium*). Univ. Calif. Pub. in Botany 30 (5): 337-436.
- LEONARD, O. A., and W. A. HARVEY
1965. Chemical control of woody plants. Calif. Agr. Exp. Sta. Bul. 812.
- LEONARD, O. A., R. K. GLENN, and D. E. BAYER
1965. Studies on the cut-surface method. I. Translocation in blue oak and madrone. Weeds 13 (4): 346-51.
- MILLER, EDWIN C.
1938. Plant physiology. New York and London: McGraw-Hill Co., Inc. 895-97.
- MOSS, VIRGIL D.
1960. Antibiotic treatment of western white pine infected with blister rust. Jour. Forestry 58 (9): 691-95.
- NELSON, C. D.
1964. The production and translocation of photosynthate-C¹⁴ in conifers. In: The formation of wood in forest trees, Martin W. Zimmermann, ed. New York and London: Academic Press. 243-57.

OFFORD, H. R.

1960. Chemical control of mistletoes. Weed Society of America Abstracts. 26-27.

PARMETER, J. R., JR., and ROBERT F. SCHARPF

1963. Dwarfmistletoe on red fir and white fir. Jour. Forestry 61 (5): 371-74.

QUICK, CLARENCE R.

1964. Experimental herbicidal control of dwarf mistletoe in some California conifers. U. S. Forest Service, Pacific Southwest Forest and Range Exp. Sta. Research Note 47. 9 pp.

REDISKE, J. H., and K. R. SHEA

1961. The production and translocation of photosynthate in dwarf mistletoe and lodgepole pine. Amer. Jour. Bot. 48 (6): 447-52.

SCHARPF, ROBERT F.

1962a. Epidemiology and parasitism of the dwarf mistletoe (*Arceuthobium campylopodium* Engelm.) in California. Ph.D. dissertation, University of California, Berkeley.

1962b. Growth rate of the endophytic system of the dwarf mistletoe on digger pine. Pacific Southwest Forest and Range Exp. Sta. Research Note 193: 1-5.

SRIVASTAVA, L. M., and K. ESAU

1961. Relation of dwarf mistletoe (*Arceuthobium*) to the xylem tissue of conifers. I. Anatomy of parasite sinkers and their connection with host xylem. Amer. Jour. Bot. 48 (2): 159-67.

VICHE, HENRY J., VIRGIL D. MOSS, and HOMER J. HARTMANN

1962. Developments in aerial application of antibiotics to control blister rust on western white pine. Jour. Forestry 60 (11): 782-84.

VITÉ, J. P., and J. A. RUDINSKY

1959. The water-conducting systems in conifers and their importance to the distribution of trunk injected chemicals. Contrib. Boyce Thompson Inst. 20 (1): 27-38.

WAGENER, WILLIS W.

1925. Mistletoe in the lower bole of incense cedar. Phytopathology 15 (10): 614-16.

WEIR, J. R.

1916. Mistletoe injury to conifers in the Northwest. U. S. Dept. Agr. Bul. 360. 39 pp.

YAMAGUCHI, S., and A. S. CRAFTS

1958. Autoradiographic method for studying absorption and translocation of herbicides using C¹⁴-labeled compounds. Hilgardia 28 (6): 161-91.

To simplify the information, it is sometimes necessary to use trade names of products or equipment. No endorsement of named products is intended nor is criticism implied of similar products not mentioned.

The journal HILGARDIA is published at irregular intervals, in volumes of about 650 to 700 pages. The number of issues per volume varies.

Single copies of any issue may be obtained free, as long as the supply lasts; please request by volume and issue number from:

**Agricultural Publications
University Hall
University of California
Berkeley, California 94720**

The limit to nonresidents of California is 10 separate titles. The limit to California residents is 20 separate titles.

The journal will be sent regularly to libraries, schools, or institutions in one of the following ways:

- 1. In exchange for similar published material on research.**
- 2. As a gift to qualified repository libraries only.**
- 3. On a subscription basis—\$7.50 a year paid in advance. All subscriptions will be started with the first number issued during a calendar year. Subscribers starting during any given year will be sent back numbers to the first of that year and will be billed for the ensuing year the following January. Make checks or money orders payable to The Regents of The University of California; send payment with order to Agricultural Publications at above address.**