
Developing a

MEAT INDEX FOR BEEF CATTLE



The idea for the meat index had its genesis in research with double-muscled animals. A marked association of rump shape and intermuscular creases with leanness and heavy musculature of the carcass was found. Double-muscled animals typically score "1" on the rump gage, whereas normal animals tend to fall in the range from 3 to 4.

Through the use of statistical techniques, the crease scores and the rump gage score can be linearly combined to allow maximum correlation of the meat index (MI) with the amount of nonfat in the carcass, expressed as a precentage of live slaughter weight.

Formulas


Based on the beef animals tested to date, the formulas developed are: MI = CS - .66 CM + 2.06 RS (for bulls); and MI = CS - .76 CM + 1.76 RS (for steers)—where CS = visibility score of the crease in the standing animal, CM = visibility score of the crease in the mov-



RUMP GAGE dimensions, exclusive of handle, are $8\frac{1}{4} \times 3\frac{1}{4}$ inches. Material is clear plexiglass.

VISIBILITY SCORING OF THE CREASE BETWEEN
THE SEMIMEMBRANOSUS AND THE
SEMITENDINOSUS MUSCLES IN BEEF ANIMALS

Visibility	Animal standing	Animal moving
	CS	CW
Marked: crease is markedly noticeable even to one unfamiliar with cattle.	1	1
Definite: crease can be seen easily by one familiar with the study, but is not marked,	2	2
Slight: crease can be seen only by the careful observance of one familiar with the study.	3	3
None: no crease is visible,	4	4



The crease (C) is between the semitendinosus (1) and the semimembranosus (2) rump muscles.

Sighting through rump gage to match one of four outlines to the aminal's rump shope.



ing animal, and RS = rump gage score. For example, if a steer had a CS score of 4, a CM score of 3, and an RS score of 3, its meat index would be $1 \times 4 - (.76 \times 3) + (1.76 \times 3) = 7.00$. The lower the MI score, the meatier is the expected carcass.

In two tests to date—one involving 13 bulls, the other 12 steers, fed out to low choice grade—the meat index was found to be highly correlated at .8 and .7 respectively (considering 1.0 as perfect prediction and 0 as no predictability), with the amount of nonfat in the carcass, expressed as a percentage of live slaughter weight. The corresponding correlations using the U.C. conformation grade were .3 and .2.

The animals used in constructing the meat indexes given here were considered normal on the basis of their individual appearance and pedigree. They were of Angus, Hereford, and Shorthorn breeding.

University of California Division of Agricultural Sciences, Agricultural Experiment Station, Berkeley, California 94720

C. 7. Felly

Director

PUBLICATION

California Agriculture

Permit No. 1127

Development of the meat index is continuing. Additional tests are to be made this year. Other intermuscular creases may also be of use to increase the accuracy of the meat index.

Wade C. Rollins is Professor and Animal Husbandman in the Experiment Sta-

tion, University of California, Davis. Collaborators in the double-muscle experiment included F. D. Carroll, Department of Animal Husbandry, and L. M. Julian, Department of Anatomy. Mrs. Moira Tanaka, Laboratory Technician, assisted with the development of the meat index.

DONATIONS FOR AGRICULTURAL RESEARCH

Contributions to the University of California, Division of Agricultural Sciences

RIVERSIDE		
Abbott Laboratories		
Allied Chemical Company		
American Potash Institute, Inc. 1,000.00 In support of research on citrus to improve production and quality		
California Fertilizer Association		
Chemagro Corporation		
Chevron Chemical Company For Plant Pathology Department work on vegetables and small fruit disease control\$ 1,000.00 For Entomology Department determinations on interactions of soils and pesticides and their implications 10,000.00		
Coachella Valley Mosquito Abatement District 12,500.00 For studies on the attractants and repellants of eye gnats by the Entomology Department		
The Dow Chemical Company		
Golf Course Superintendents Association of America 1,500.00 For turfgrass research by Agronomy Department		
Hazleton Laboratories, Incorporated		
Stauffer Chemical CompanyBye-Hoe herbicide incorporator unit and trailer		

For use by Department of Vegetable Crops

Thompson-Hayward Chemical Company\$ In support of weed control research program of the Department of Horticultural Science	500.00
Ventura County Citrus Growers Committee, Inc For use by Horticultural Science Department on fruit harvesting research	500.00
DAVIS	
Beken Bros	1,650.00
California Sugar Beet Processors American Crystal Sugar Company Holly Sugar Corporation Spreckels Sugar Company Union Sugar Various Donors	
California Beet Growers Association, Ltd	11,000.00
Chevron Chemical Company To continue support of research in stone fruit diseases— Plant Pathology In support of research on range improvement— Agronomy	
Mr. Lloyd Fox One hundred mated que To Department of Entomology for research purposes	een bees
General Electric Company	500.00
Hewlett-Packard Company Scaler Timer, High Voltage Power Supply Scintillation For research in soil water movement—Water Science and Engineering	Detector
Humboldt County Wool Growers' Association\$ To be added to Various Donor Fund in Sheep Foot Rot Research	100.00