

New Mite Predators

four species from Guatemala show promise in southern California

C. A. Fleschner

The Guatemalan *Stethorus*—a small, black, lady beetle mite predator—is being propagated by the thousand in the insectary at Riverside for release in southern California avocado and citrus groves.

As a result of an exploratory trip through Mexico and Central America during the winter of 1953-54 to study the natural balance of pests of avocado trees in their native habitat, several species of natural enemies controlling mite pests of wild avocado trees were found in the Guatemalan highlands. Although Guatemala is a tropical country,

frost may occur in the highlands above 5,000' during the colder months, so it was thought that beneficial species collected from this area could possibly be established in southern California.

Consequently, after methods were developed for their propagation, nine species of these mite predators were shipped to Riverside in the spring of 1955, and four of these—one cheyletid mite, two typhlodromid mites, and one *Stethorus*—are being propagated. All four are new, undescribed species.

Stethorus, particularly, shows promise. Over 7,000 of this predator have already

been released in the southern counties where it is working and reproducing well, although it remains to be seen how satisfactorily it will work through the winter months. Large numbers will be used experimentally in mass liberations at critical times in attempts to prevent the development of injurious plant-feeding mite infestations.

Since *Stethorus* feeds on a wide range of mite species, it may be used in this experimental manner not only on avocado and citrus but on various truck and field crops and other trees. This is also true of the cheyletid mite and the two typhlodromid mites, which—as laboratory studies have shown—will feed on all mite pests of avocado and citrus trees except the citrus bud mite.

Propagation of these beneficial species will be continued until they have been given an opportunity to become established in southern California groves.

C. A. Fleschner is Associate Entomologist, University of California, Riverside.

The above progress report is based on Research Project No. 1495.

FRUIT SIZE

Continued from preceding page

from each culture were tested in duplicate determinations for their content of calcium, potassium, magnesium, and sodium, the results of which are shown in the larger table. The calcium percentage in the peel was approximately doubled while that of potassium was more than tripled in the extreme. Hoagland's nutrient solution regularly contains 159 ppm of calcium. When the concentration of calcium in the nutrient solution was increased above this strength—as shown in the larger table—the calcium percentage in the peel made its greatest increase. Above a concentration of 237 ppm of calcium in the nutrient, calcium percentage in the peel increased only slightly.

The peel readily absorbed large amounts of potassium when the concentration of potassium in the nutrient solution was increased and that of calcium was simultaneously decreased. In the

Yields and Sizes of Lemon Fruits in the Silver Stage of Maturity Grown on Trees in Soil Cultures That Received Similar Nutrient Solutions Except for the Ratio of Calcium and Potassium

Culture	Calcium	Potassium	No. of fruit	Wt. of fruit grms.	Diameter of fruit ins.
1	318	0	12	1134	2.095
2	298	39	23	2335	2.165
3	277	77	15	2400	2.169
4	237	155	13	1503	2.193
5	159	310	18	2027	2.228
6	80	467	19	2227	2.252
7	40	545	11	1299	2.272
8	0	622	8	1100	2.374

peel there were very small increases in the percentages of magnesium accompanying the increased calcium and reduced potassium percentages though the magnesium concentrations in the nutrient solutions were similar for all cultures.

In the pulp the calcium percentages, while relatively low in comparison with those of potassium, showed a consistent

increase, whereas the magnesium—constant in each nutrient—showed minor increases as calcium in the nutrient was increased and potassium decreased.

Lemon fruit therefore responded very well to changes in the calcium-potassium content in the nutrient solution applied to the soil when the concentration of the applied other nutrient elements was not varied in any of the cultures. The larger table also shows that the calcium and potassium percentages in the dry matter of lemon flowers can be markedly affected by varying the ratio of calcium to potassium in the nutrient applied to soil cultures. The high calcium content of many irrigation waters should make the potassium supply of considerable importance.

A. R. C. Haas is Plant Physiologist, Emeritus, University of California, Riverside.

Joseph N. Brusca is Principal Laboratory Technician, University of California, Riverside.

The above progress report is based on Research Project No. 1086.

CITRUS

Continued from page 12

tion of importation bud of wood or cuttings was modified, and introduction of citrus budwood from other countries is permissible under quarantine measures.

Quarantine regulations—imposed in the 1930's—prohibited importation of budwood or cuttings from out of state because of the danger of introducing contagious diseases with the incoming material.

Material introduced under current quarantine measures is carefully screened and indexed for various diseases at the U.S.D.A. Plant Industry Station greenhouses at Beltsville, Maryland. Satisfactory material can be released to California where it will again be placed in isolation quarantine greenhouses. The introductions will be budded into various rootstock combinations and into different seedlings to index for miscellaneous virus and other diseases. Material found to be satisfactory will be released

from quarantine to governmental agencies for experimental field trials.

The variety orchard provides a wide source of propagating material of miscellaneous varieties and a varied population of genes for breeding studies. Wide use of such trees is made in rootstock tests and in the search for nematode and disease-resistant stocks.

W. P. Bitters is Associate Horticulturist, University of California, Riverside.

The above progress report is based on Research Project No. 194.